

Discrete Mathematics & Computational Structures  
Lattice-Point Counting in Convex Polytopes  
(2) A garely of discrete volumes

Yoshio Okamoto

Tokyo Institute of Technology

April 30, 2009

"Last updated: 2009/05/01 15:49"

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes

# Convex polytopes in dimension 1

## Convex polytopes in dimension 1

= straight line segments

- Integral segment  $[a, b]$ ,  $a, b \in \mathbb{Z}$ ,  $a < b$

$$\#([a, b] \cap \mathbb{Z}) = b - a + 1$$



## Convex polytopes in dimension 1

## Convex polytopes in dimension 1

= straight line segments

- Integral segment  $[a, b]$ ,  $a, b \in \mathbb{Z}$ ,  $a < b$

$$\#([a, b] \cap \mathbb{Z}) = b - a + 1$$



- Rational segment  $[a/b, c/d]$ ,  $a, b, c, d \in \mathbb{Z}$ ,  $a/b < c/d$


$$\#([a/b, c/d] \cap \mathbb{Z}) = \lfloor c/d \rfloor - \lfloor (a-1)/b \rfloor$$

(Exercise 2.1)

# Convex polytopes in dimension 2

Convex polytopes in dimension 2

= convex polygons




Lattice-point counting is a topic of Sect. 2.6 and 2.7

Convex polytopes in dimension  $d$ Convex polytopes in dimension  $d$ 

= convex hulls of a finite set of points

For  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\} \subset \mathbb{R}^d$ 

$$\mathcal{P} = \left\{ \lambda_1 \mathbf{v}_1 + \lambda_2 \mathbf{v}_2 + \dots + \lambda_n \mathbf{v}_n : \begin{array}{l} \text{all } \lambda_k \geq 0 \text{ and} \\ \lambda_1 + \lambda_2 + \dots + \lambda_n = 1 \end{array} \right\},$$

that is, the smallest convex set containing  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ 


Convex polytopes in dimension  $d$ , another point of view

Convex polytopes in dimension  $d$

= bounded intersections of finitely many half-spaces

For  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m \in \mathbb{R}^d$  and  $b_1, b_2, \dots, b_m \in \mathbb{R}$

$$\mathcal{P} = \{\mathbf{x} : \mathbf{a}_i \cdot \mathbf{x} \leq b_i \text{ for all } i = 1, 2, \dots, m\}$$



# Convex polytopes

Convex polytopes are

- convex hulls of a finite set of points  
(**vertex description**, v-polytopes)

or

- bounded intersections of finitely many half-spaces  
(**hyperplane description**, h-polytopes)

# Convex polytopes

Convex polytopes are

- convex hulls of a finite set of points  
(**vertex description**, v-polytopes)

or

- bounded intersections of finitely many half-spaces  
(**hyperplane description**, h-polytopes)

Theorem (Main theorem for polytopes, Minkowski-Weyl Theorem)

- Every v-polytope is an h-polytope
- Every h-polytope is a v-polytope

Proof: See Appendix A in the textbook

# Dimension of convex polytopes

$\mathcal{P}$  a convex polytope

## Definition (Dimension)

The **dimension** of  $\mathcal{P}$  is the dimension of the span of  $\mathcal{P}$ , where

$$\text{span } \mathcal{P} := \{\mathbf{x} + \lambda(\mathbf{y} - \mathbf{x}) : \mathbf{x}, \mathbf{y} \in \mathcal{P}, \lambda \in \mathbb{R}\}$$

## Definition ( $d$ -Polytope)

If the dimension of  $\mathcal{P}$  is  $d$ , then we write


- $\dim \mathcal{P} = d$
- $\mathcal{P}$  is a  **$d$ -polytope**

# Valid inequalities

$\mathcal{P} \subseteq \mathbb{R}^d$  a convex polytope;  $\mathbf{a} \in \mathbb{R}^d$ ,  $b \in \mathbb{R}$

## Definition (Valid inequality)

The inequality  $\mathbf{a} \cdot \mathbf{x} \leq b$  is a **valid inequality** for  $\mathcal{P}$  if  
 $\mathbf{a} \cdot \mathbf{z} \leq b$  for all  $\mathbf{z} \in \mathcal{P}$




## Faces of a convex polytope

$\mathcal{P} \subseteq \mathbb{R}^d$  a convex polytope

### Definition (Face)

$\mathcal{F}$  is a **face** of  $\mathcal{P}$  if  $\exists$  a valid inequality  $\mathbf{a} \cdot \mathbf{x} \leq b$  for  $\mathcal{P}$  s.t.

$$\mathcal{F} = \mathcal{P} \cap \{\mathbf{x} : \mathbf{a} \cdot \mathbf{x} = b\}$$



### Remark

- Every face of a convex polytope is also a convex polytope
- $\mathcal{P}$  and  $\emptyset$  are faces of  $\mathcal{P}$

# Names of faces


$\mathcal{P} \subseteq \mathbb{R}^d$  a convex polytope

## Definition

0-dimensional face **vertex** of  $\mathcal{P}$

1-dimensional face **edge** of  $\mathcal{P}$

$(d-1)$ -dimensional face **facet** of  $\mathcal{P}$





# Simplices

## Fact

Every convex  $d$ -polytope has at least  $d+1$  vertices

## Definition (Simplex)

A convex  $d$ -polytope is a  **$d$ -simplex** if it has exactly  $d+1$  vertices



# Integral polytopes and rational polytopes

## Definition (Integral polytope)

A convex polytope is **integral** if all of its vertices have integer coordinates

## Definition (Rational polytope)

A convex polytope is **rational** if all of its vertices have rational coordinates

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes

# Unit cubes


## Definition (Unit $d$ -cube $\square$ )

- Vertices

$$\{(x_1, x_2, \dots, x_d) : \text{all } x_k = 0 \text{ or } 1\}$$


- Hyperplane description

$$\square = \{(x_1, x_2, \dots, x_d) \in \mathbb{R}^d : 0 \leq x_k \leq 1 \text{ for all } k\}.$$



## Question

What's the number of integer points in  $t \square$  ( $t \in \mathbb{Z}_{>0}$ )?



$$\#(t \square \cap \mathbb{Z}^d) = \#([0, t]^d \cap \mathbb{Z}^d) = (t + 1)^d$$

## Lattice-point enumerators

$\mathcal{P} \subseteq \mathbb{R}^d$  not necessarily a convex polytope

Definition (Lattice-point enumerator)

The **lattice-point enumerator** for  $t\mathcal{P}$  is defined as

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^d)$$

### Remarks

- $L_{\mathcal{P}}(t) = \#(\mathcal{P} \cap \frac{1}{t}\mathbb{Z}^d)$

## Lattice-point enumerators

$\mathcal{P} \subseteq \mathbb{R}^d$  not necessarily a convex polytope

Definition (Lattice-point enumerator)

The **lattice-point enumerator** for  $t\mathcal{P}$  is defined as

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^d)$$

### Remarks

- $L_{\mathcal{P}}(t) = \#(\mathcal{P} \cap \frac{1}{t}\mathbb{Z}^d)$
- $L_{\square}(t) = (t + 1)^d$

$(t + 1)^d$  is the generating function of ...

$$(t + 1)^d = \sum_{k=0}^d \binom{d}{k} t^k,$$

where  $\binom{d}{k}$  is a **binomial coefficient** defined as follows


### Definition (Binomial coefficient)

For  $m \in \mathbb{C}, n \in \mathbb{Z}_{>0}$

$$\binom{m}{n} := \frac{m(m-1)(m-2) \cdots (m-n+1)}{n!} \quad (1)$$

# What about the interior?

$$L_{\square^\circ}(t) = \#(t \square^\circ \cap \mathbb{Z}^d) = \#((0, t)^d \cap \mathbb{Z}^d) = (t - 1)^d$$



**Remark**

$$L_{\square^\circ}(t) = (-1)^d L_{\square}(-t)$$

# Ehrhart series — another tool for studying discrete volume

$$\mathcal{P} \subseteq \mathbb{R}^d$$

## Definition (Ehrhart series)

The **Ehrhart series** of  $\mathcal{P}$  is the generating fn of  $L_{\mathcal{P}}(t)$ :

$$\text{Ehr}_{\mathcal{P}}(z) := 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t$$

Namely,

$$\text{Ehr}_{\square}(z) = 1 + \sum_{t \geq 1} (t+1)^d z^t$$

# Eulerian numbers

## Definition (Eulerian number)

For  $0 \leq k \leq d$ , the **Eulerian number**  $A(d, k)$  is defined through

$$\sum_{j \geq 0} j^d z^j = \frac{\sum_{k=0}^d A(d, k) z^k}{(1-z)^{d+1}} \quad (2)$$

Then,

$$\text{Ehr}_{\square}(z) = 1 + \sum_{t \geq 1} (t+1)^d z^t$$

# Eulerian numbers

## Definition (Eulerian number)

For  $0 \leq k \leq d$ , the **Eulerian number**  $A(d, k)$  is defined through

$$\sum_{j \geq 0} j^d z^j = \frac{\sum_{k=0}^d A(d, k) z^k}{(1 - z)^{d+1}} \quad (2)$$

Then,

$$\text{Ehr}_{\square}(z) = 1 + \sum_{t \geq 1} (t + 1)^d z^t = \sum_{t \geq 0} (t + 1)^d z^t$$

# Eulerian numbers

## Definition (Eulerian number)

For  $0 \leq k \leq d$ , the **Eulerian number**  $A(d, k)$  is defined through

$$\sum_{j \geq 0} j^d z^j = \frac{\sum_{k=0}^d A(d, k) z^k}{(1-z)^{d+1}} \quad (2)$$

Then,

$$\text{Ehr}_{\square}(z) = 1 + \sum_{t \geq 1} (t+1)^d z^t = \sum_{t \geq 0} (t+1)^d z^t = \frac{1}{z} \sum_{t \geq 1} t^d z^t$$

# Eulerian numbers

## Definition (Eulerian number)

For  $0 \leq k \leq d$ , the **Eulerian number**  $A(d, k)$  is defined through

$$\sum_{j \geq 0} j^d z^j = \frac{\sum_{k=0}^d A(d, k) z^k}{(1 - z)^{d+1}} \quad (2)$$

Then,

$$\begin{aligned} \text{Ehr}_{\square}(z) &= 1 + \sum_{t \geq 1} (t + 1)^d z^t = \sum_{t \geq 0} (t + 1)^d z^t = \frac{1}{z} \sum_{t \geq 1} t^d z^t \\ &= \frac{\sum_{k=1}^d A(d, k) z^{k-1}}{(1 - z)^{d+1}} \end{aligned}$$

# What's an Eulerian number?

## Fact

$A(d, k) = \# \text{ permutations of } \{1, \dots, d\} \text{ with } k-1 \text{ ascents}$

$$d = 6 : 1 \quad 4 \quad 2 \quad 5 \quad 6 \quad 3$$

# What's an Eulerian number?

## Fact

$A(d, k) = \# \text{ permutations of } \{1, \dots, d\} \text{ with } k-1 \text{ ascents}$

$$d = 6 \quad : 1 \quad 4 \mid 2 \quad 5 \quad 6 \mid 3$$

# What's an Eulerian number?

## Fact

$A(d, k) = \# \text{ permutations of } \{1, \dots, d\} \text{ with } k-1 \text{ ascents}$

$$d = 6, k = 3: 1 \quad 4 \mid 2 \quad 5 \quad 6 \mid 3$$

## What's an Eulerian number?

## Fact

$$A(d, k) = \# \text{ permutations of } \{1, \dots, d\} \text{ with } k-1 \text{ ascents}$$

$$d = 6, k = 3: 1 \quad 4 \mid 2 \quad 5 \quad 6 \mid 3$$

## Properties (Exercise 2.8)

$$1 \leq k \leq d$$

$$A(d, k) = A(d, d + 1 - k),$$

$$A(d, k) = (d - k + 1) A(d - 1, k - 1) + k A(d - 1, k),$$

$$\sum_{k=0}^d A(d, k) = d!, \tag{3}$$

$$A(d, k) = \sum_{j=0}^k (-1)^j \binom{d+1}{j} (k-j)^d$$

## Summary: The unit cube

### Theorem 2.1

- (a) The lattice-point enumerator of  $\square$  is the polynomial

$$L_{\square}(t) = (t + 1)^d = \sum_{k=0}^d \binom{d}{k} t^k$$

- (b) Its evaluation at negative integers yields the relation

$$(-1)^d L_{\square}(-t) = L_{\square^\circ}(t)$$

- (c) The Ehrhart series of  $\square$  is  $\text{Ehr}_{\square}(z) = \frac{\sum_{k=1}^d A(d, k) z^{k-1}}{(1 - z)^{d+1}}$

□

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes


# Standard simplices

## Definition (Standard $d$ -simplex)

- Vertices  
 $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_d$  and  $\mathbf{0}$ , where  
 $\mathbf{e}_j$  is the unit vector  $(0, \dots, 1, \dots, 0)$  with a 1 in the  $j$ -th position
- Hyperplane description

$$\Delta = \left\{ (x_1, x_2, \dots, x_d) \in \mathbb{R}^d : \begin{array}{l} x_1 + x_2 + \cdots + x_d \leq 1, \\ \text{all } x_k \geq 0 \end{array} \right\}$$

## Example: Standard 3-simplex



The dilated standard simplex  $t\Delta$ 

$$t\Delta = \left\{ (x_1, x_2, \dots, x_d) \in \mathbb{R}^d : \begin{array}{l} x_1 + x_2 + \cdots + x_d \leq t, \\ \text{all } x_k \geq 0 \end{array} \right\}$$

Let's compute the discrete volume and the Ehrhard series!

We need a trick

- $\Delta$  involves an **inequality**
- The example from Lecture 1 involves **equalities only**

# The dilated standard simplex $t\Delta$

$$t\Delta = \left\{ (x_1, x_2, \dots, x_d) \in \mathbb{R}^d : \begin{array}{l} x_1 + x_2 + \cdots + x_d \leq t, \\ \text{all } x_k \geq 0 \end{array} \right\}$$

Let's compute the discrete volume and the Ehrhard series!

We need a trick

- $\Delta$  involves an **inequality**
- The example from Lecture 1 involves **equalities only**

Trick  $\rightarrow$   $\left\{ \begin{array}{l} \text{Transform an inequality to equalities by introducing} \\ \text{an extra coordinate} \end{array} \right.$

## Slack variables

- Want to count all integer solutions  $(m_1, m_2, \dots, m_d) \in \mathbb{Z}_{\geq 0}^d$  to

$$m_1 + m_2 + \cdots + m_d \leq t \quad (4)$$

## Slack variables

- Want to count all integer solutions  $(m_1, m_2, \dots, m_d) \in \mathbb{Z}_{\geq 0}^d$  to

$$m_1 + m_2 + \cdots + m_d \leq t \quad (4)$$

- Let  $m_{d+1} = \text{RHS} - \text{LHS} \geq 0$

Such a variable  $m_{d+1}$  is called a **slack variable**

## Slack variables

- Want to count all integer solutions  $(m_1, m_2, \dots, m_d) \in \mathbb{Z}_{\geq 0}^d$  to

$$m_1 + m_2 + \cdots + m_d \leq t \quad (4)$$

- Let  $m_{d+1} = \text{RHS} - \text{LHS} \geq 0$
- Then

# integer solutions  $(m_1, m_2, \dots, m_d) \in \mathbb{Z}_{\geq 0}^d$  to  
 $m_1 + m_2 + \cdots + m_d \leq t$

||

# integer solutions  $(m_1, m_2, \dots, m_{d+1}) \in \mathbb{Z}_{\geq 0}^{d+1}$  to  
 $m_1 + m_2 + \cdots + m_{d+1} = t$

Such a variable  $m_{d+1}$  is called a **slack variable**

## Discrete volume of a standard simplex

- Similarly to Lecture 1

$$\#(t\Delta \cap \mathbb{Z}^d)$$

$$= \text{const} \left( \left( \sum_{m_1 \geq 0} z^{m_1} \right) \left( \sum_{m_2 \geq 0} z^{m_2} \right) \cdots \left( \sum_{m_{d+1} \geq 0} z^{m_{d+1}} \right) z^{-t} \right)$$

(5)

## Discrete volume of a standard simplex

- Similarly to Lecture 1

$$\begin{aligned}\#(t\Delta \cap \mathbb{Z}^d) &= \text{const} \left( \left( \sum_{m_1 \geq 0} z^{m_1} \right) \left( \sum_{m_2 \geq 0} z^{m_2} \right) \cdots \left( \sum_{m_{d+1} \geq 0} z^{m_{d+1}} \right) z^{-t} \right) \\ &= \text{const} \left( \frac{1}{(1-z)^{d+1} z^t} \right) \quad (5)\end{aligned}$$

## Discrete volume of a standard simplex

- Similarly to Lecture 1

$$\begin{aligned}
 & \#(t\Delta \cap \mathbb{Z}^d) \\
 &= \text{const} \left( \left( \sum_{m_1 \geq 0} z^{m_1} \right) \left( \sum_{m_2 \geq 0} z^{m_2} \right) \cdots \left( \sum_{m_{d+1} \geq 0} z^{m_{d+1}} \right) z^{-t} \right) \\
 &= \text{const} \left( \frac{1}{(1-z)^{d+1} z^t} \right)
 \end{aligned} \tag{5}$$

- Now use the **binomial series**

$$\frac{1}{(1-z)^{d+1}} = \sum_{k \geq 0} \binom{d+k}{d} z^k \quad \text{for } d \geq 0 \tag{6}$$

## Discrete volume of a standard simplex

- Similarly to Lecture 1

$$\begin{aligned}
 & \#(t\Delta \cap \mathbb{Z}^d) \\
 &= \text{const} \left( \left( \sum_{m_1 \geq 0} z^{m_1} \right) \left( \sum_{m_2 \geq 0} z^{m_2} \right) \cdots \left( \sum_{m_{d+1} \geq 0} z^{m_{d+1}} \right) z^{-t} \right) \\
 &= \text{const} \left( \frac{1}{(1-z)^{d+1} z^t} \right)
 \end{aligned} \tag{5}$$

- Now use the **binomial series**

$$\frac{1}{(1-z)^{d+1}} = \sum_{k \geq 0} \binom{d+k}{d} z^k \quad \text{for } d \geq 0 \tag{6}$$

- That gives  $L_\Delta(t) := \#(t\Delta \cap \mathbb{Z}^d) = \binom{d+t}{d}$

What about the interior  $\Delta^\circ$ ?

$$L_{\Delta^\circ}(t)$$

$$= \# \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}_{>0}^d : m_1 + m_2 + \cdots + m_d < t \right\}$$

What about the interior  $\Delta^\circ$ ?

$$L_{\Delta^\circ}(t)$$

$$= \# \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}_{>0}^d : m_1 + m_2 + \dots + m_d < t \right\}$$

$$= \# \left\{ (m_1, m_2, \dots, m_{d+1}) \in \mathbb{Z}_{>0}^{d+1} : m_1 + m_2 + \dots + m_{d+1} = t \right\}$$

What about the interior  $\Delta^\circ$ ?

$$L_{\Delta^\circ}(t)$$

$$= \# \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}_{>0}^d : m_1 + m_2 + \dots + m_d < t \right\}$$

$$= \# \left\{ (m_1, m_2, \dots, m_{d+1}) \in \mathbb{Z}_{>0}^{d+1} : m_1 + m_2 + \dots + m_{d+1} = t \right\}$$

$$= \text{const} \left( \left( \sum_{m_1>0} z^{m_1} \right) \left( \sum_{m_2>0} z^{m_2} \right) \cdots \left( \sum_{m_{d+1}>0} z^{m_{d+1}} \right) z^{-t} \right)$$

What about the interior  $\Delta^\circ$ ?

$$L_{\Delta^\circ}(t)$$

$$\begin{aligned} &= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}_{>0}^d : m_1 + m_2 + \dots + m_d < t\} \\ &= \#\{(m_1, m_2, \dots, m_{d+1}) \in \mathbb{Z}_{>0}^{d+1} : m_1 + m_2 + \dots + m_{d+1} = t\} \\ &= \text{const} \left( \left( \sum_{m_1>0} z^{m_1} \right) \left( \sum_{m_2>0} z^{m_2} \right) \cdots \left( \sum_{m_{d+1}>0} z^{m_{d+1}} \right) z^{-t} \right) \\ &= \text{const} \left( \left( \frac{z}{1-z} \right)^{d+1} z^{-t} \right) \end{aligned}$$

What about the interior  $\Delta^\circ$ ?

$$L_{\Delta^\circ}(t)$$

$$\begin{aligned}
 &= \# \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}_{>0}^d : m_1 + m_2 + \cdots + m_d < t \right\} \\
 &= \# \left\{ (m_1, m_2, \dots, m_{d+1}) \in \mathbb{Z}_{>0}^{d+1} : m_1 + m_2 + \cdots + m_{d+1} = t \right\} \\
 &= \text{const} \left( \left( \sum_{m_1>0} z^{m_1} \right) \left( \sum_{m_2>0} z^{m_2} \right) \cdots \left( \sum_{m_{d+1}>0} z^{m_{d+1}} \right) z^{-t} \right) \\
 &= \text{const} \left( \left( \frac{z}{1-z} \right)^{d+1} z^{-t} \right) \\
 &= \text{const} \left( z^{d+1-t} \sum_{k \geq 0} \binom{d+k}{d} z^k \right)
 \end{aligned}$$

What about the interior  $\Delta^\circ$ ?

$$L_{\Delta^\circ}(t)$$

$$\begin{aligned}
 &= \# \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}_{>0}^d : m_1 + m_2 + \cdots + m_d < t \right\} \\
 &= \# \left\{ (m_1, m_2, \dots, m_{d+1}) \in \mathbb{Z}_{>0}^{d+1} : m_1 + m_2 + \cdots + m_{d+1} = t \right\} \\
 &= \text{const} \left( \left( \sum_{m_1>0} z^{m_1} \right) \left( \sum_{m_2>0} z^{m_2} \right) \cdots \left( \sum_{m_{d+1}>0} z^{m_{d+1}} \right) z^{-t} \right) \\
 &= \text{const} \left( \left( \frac{z}{1-z} \right)^{d+1} z^{-t} \right) \\
 &= \text{const} \left( z^{d+1-t} \sum_{k \geq 0} \binom{d+k}{d} z^k \right) \\
 &= \binom{t-1}{d}
 \end{aligned}$$

What about the interior  $\Delta^\circ$ ?

$$L_{\Delta^\circ}(t)$$

$$\begin{aligned}
 &= \# \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}_{>0}^d : m_1 + m_2 + \cdots + m_d < t \right\} \\
 &= \# \left\{ (m_1, m_2, \dots, m_{d+1}) \in \mathbb{Z}_{>0}^{d+1} : m_1 + m_2 + \cdots + m_{d+1} = t \right\} \\
 &= \text{const} \left( \left( \sum_{m_1>0} z^{m_1} \right) \left( \sum_{m_2>0} z^{m_2} \right) \cdots \left( \sum_{m_{d+1}>0} z^{m_{d+1}} \right) z^{-t} \right) \\
 &= \text{const} \left( \left( \frac{z}{1-z} \right)^{d+1} z^{-t} \right) \\
 &= \text{const} \left( z^{d+1-t} \sum_{k \geq 0} \binom{d+k}{d} z^k \right) \\
 &= \binom{t-1}{d} \stackrel{\text{Ex. 2.10}}{=} (-1)^d \binom{d-t}{d}
 \end{aligned}$$

## Summary: The standard $d$ -simplex

### Theorem 2.2

- (a) The lattice-point enumerator of  $\Delta$  is the polynomial

$$L_{\Delta}(t) = \binom{d+t}{d}$$


- (b) Its evaluation at negative integers yields

$$(-1)^d L_{\Delta}(-t) = L_{\Delta^{\circ}}(t)$$

- (c) The Ehrhart series of  $\Delta$  is  $\text{Ehr}_{\Delta}(z) = \frac{1}{(1-z)^{d+1}}$

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes

## Example: The 3-dimensional pyramid



# Pyramids

## Definition ( $d$ -Dimensional pyramid)

- Vertices

$$\{(x_1, x_2, \dots, x_{d-1}, 0) : \text{all } x_k = 0 \text{ or } 1\} \cup \{(0, 0, \dots, 0, 1)\}$$

- Hyperplane description

$$\mathcal{P} = \left\{ (x_1, x_2, \dots, x_d) \in \mathbb{R}^d : \begin{array}{l} 0 \leq x_k \leq 1 - x_d \leq 1 \\ \text{for all } k = 1, \dots, d-1 \end{array} \right\} \quad (9)$$

## Remark

$d$ -dim. pyramid  $\subseteq$   $d$ -dim. unit cube

## Lattice-point enumerator of a pyramid

$$\begin{aligned} L_{\mathcal{P}}(t) &= \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : \begin{array}{l} 0 \leq m_k \leq t - m_d \leq t \\ \text{for all } k = 1, \dots, d-1 \end{array} \right\} \\ &= \sum_{m_d=0}^t (t - m_d + 1)^{d-1} \\ &= \sum_{k=1}^{t+1} k^{d-1} \end{aligned}$$

Question

What's the last sum?

## Bernoulli polynomials and Bernoulli numbers

### Definition (Bernoulli polynomial)

The **Bernoulli polynomials**  $B_k(x)$  are defined via the generating fn

$$\frac{ze^{xz}}{e^z - 1} = \sum_{k \geq 0} \frac{B_k(x)}{k!} z^k \quad (8)$$

First few Bernoulli polynomials:  $k = 0, 1, 2, 3, \dots$

$$1, x - \frac{1}{2}, x^2 - x + \frac{1}{6}, x^3 - \frac{3}{2}x^2 + \frac{1}{2}x, \dots$$

### Definition (Bernoulli number)

The **Bernoulli numbers** are  $B_k := B_k(0)$

# A lemma to show a connection...

## Lemma 2.3

$$\sum_{k=0}^{n-1} k^{d-1} = \frac{1}{d} (B_d(n) - B_d) \quad \text{for integers } d \geq 1, n \geq 2$$

# A lemma to show a connection...

## Lemma 2.3

$$\sum_{k=0}^{n-1} k^{d-1} = \frac{1}{d} (B_d(n) - B_d) \quad \text{for integers } d \geq 1, n \geq 2$$

Proof:

$$\sum_{d \geq 0} \frac{B_d(n) - B_d}{d!} z^d$$

# A lemma to show a connection...

## Lemma 2.3

$$\sum_{k=0}^{n-1} k^{d-1} = \frac{1}{d} (B_d(n) - B_d) \quad \text{for integers } d \geq 1, n \geq 2$$

Proof:

$$\sum_{d \geq 0} \frac{B_d(n) - B_d}{d!} z^d = z \frac{e^{nz} - 1}{e^z - 1}$$

## A lemma to show a connection...

### Lemma 2.3

$$\sum_{k=0}^{n-1} k^{d-1} = \frac{1}{d} (B_d(n) - B_d) \quad \text{for integers } d \geq 1, n \geq 2$$

Proof:

$$\sum_{d \geq 0} \frac{B_d(n) - B_d}{d!} z^d = z \frac{e^{nz} - 1}{e^z - 1} = z \sum_{k=0}^{n-1} e^{kz}$$

## A lemma to show a connection...

## Lemma 2.3

$$\sum_{k=0}^{n-1} k^{d-1} = \frac{1}{d} (B_d(n) - B_d) \quad \text{for integers } d \geq 1, n \geq 2$$

Proof:

$$\begin{aligned} \sum_{d \geq 0} \frac{B_d(n) - B_d}{d!} z^d &= z \frac{e^{nz} - 1}{e^z - 1} = z \sum_{k=0}^{n-1} e^{kz} \\ &= z \sum_{k=0}^{n-1} \sum_{j \geq 0} \frac{(kz)^j}{j!} \end{aligned}$$

## A lemma to show a connection...

## Lemma 2.3

$$\sum_{k=0}^{n-1} k^{d-1} = \frac{1}{d} (B_d(n) - B_d) \quad \text{for integers } d \geq 1, n \geq 2$$

Proof:

$$\begin{aligned} \sum_{d \geq 0} \frac{B_d(n) - B_d}{d!} z^d &= z \frac{e^{nz} - 1}{e^z - 1} = z \sum_{k=0}^{n-1} e^{kz} \\ &= z \sum_{k=0}^{n-1} \sum_{j \geq 0} \frac{(kz)^j}{j!} = \sum_{j \geq 0} \left( \sum_{k=0}^{n-1} k^j \right) \frac{z^{j+1}}{j!} \end{aligned}$$

## A lemma to show a connection...

## Lemma 2.3

$$\sum_{k=0}^{n-1} k^{d-1} = \frac{1}{d} (B_d(n) - B_d) \quad \text{for integers } d \geq 1, n \geq 2$$

Proof:

$$\begin{aligned} \sum_{d \geq 0} \frac{B_d(n) - B_d}{d!} z^d &= z \frac{e^{nz} - 1}{e^z - 1} = z \sum_{k=0}^{n-1} e^{kz} \\ &= z \sum_{k=0}^{n-1} \sum_{j \geq 0} \frac{(kz)^j}{j!} = \sum_{j \geq 0} \left( \sum_{k=0}^{n-1} k^j \right) \frac{z^{j+1}}{j!} \\ &= \sum_{j \geq 1} \left( \sum_{k=0}^{n-1} k^{j-1} \right) \frac{z^j}{(j-1)!} \end{aligned}$$

and compare the both sides

## Lattice-point enumerator of a pyramid, cont'd

$$\begin{aligned} L_{\mathcal{P}}(t) &= \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : \begin{array}{l} 0 \leq m_k \leq 1 - m_d \leq 1 \\ \text{for all } k = 1, \dots, d-1 \end{array} \right\} \\ &= \sum_{m_d=0}^t (t - m_d + 1)^{d-1} \\ &= \sum_{k=1}^{t+1} k^{d-1} \end{aligned}$$

(10)

## Lattice-point enumerator of a pyramid, cont'd

$$\begin{aligned}
 L_{\mathcal{P}}(t) &= \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : \begin{array}{l} 0 \leq m_k \leq 1 - m_d \leq 1 \\ \text{for all } k = 1, \dots, d-1 \end{array} \right\} \\
 &= \sum_{m_d=0}^t (t - m_d + 1)^{d-1} \\
 &= \sum_{k=1}^{t+1} k^{d-1} \\
 &= \sum_{k=0}^{t+1} k^{d-1} \quad (\text{if } d \geq 2)
 \end{aligned} \tag{10}$$

## Lattice-point enumerator of a pyramid, cont'd

$$\begin{aligned}
 L_{\mathcal{P}}(t) &= \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : \begin{array}{l} 0 \leq m_k \leq 1 - m_d \leq 1 \\ \text{for all } k = 1, \dots, d-1 \end{array} \right\} \\
 &= \sum_{m_d=0}^t (t - m_d + 1)^{d-1} \\
 &= \sum_{k=1}^{t+1} k^{d-1} \\
 &= \sum_{k=0}^{t+1} k^{d-1} \quad (\text{if } d \geq 2) \\
 &= \frac{1}{d} (B_d(t+2) - B_d)
 \end{aligned} \tag{10}$$

How about the interior  $\mathcal{P}^\circ$ ?

$$L_{\mathcal{P}^\circ}(t) = \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : \begin{array}{l} 0 < m_k < 1 - m_d < 1 \\ \text{for all } k = 1, \dots, d-1 \end{array} \right\}$$

How about the interior  $\mathcal{P}^\circ$ ?

$$\begin{aligned}L_{\mathcal{P}^\circ}(t) &= \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : \begin{array}{l} 0 < m_k < 1 - m_d < 1 \\ \text{for all } k = 1, \dots, d-1 \end{array} \right\} \\&= \sum_{m_d=1}^{t-1} (t - m_d - 1)^{d-1}\end{aligned}$$

How about the interior  $\mathcal{P}^\circ$ ?

$$\begin{aligned}L_{\mathcal{P}^\circ}(t) &= \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : \begin{array}{l} 0 < m_k < 1 - m_d < 1 \\ \text{for all } k = 1, \dots, d-1 \end{array} \right\} \\&= \sum_{m_d=1}^{t-1} (t - m_d - 1)^{d-1} \\&= \sum_{k=0}^{t-2} k^{d-1} = \frac{1}{d} (B_d(t-1) - B_d)\end{aligned}$$

How about the interior  $\mathcal{P}^\circ$ ?

$$\begin{aligned}
 L_{\mathcal{P}^\circ}(t) &= \left\{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : \begin{array}{l} 0 < m_k < 1 - m_d < 1 \\ \text{for all } k = 1, \dots, d-1 \end{array} \right\} \\
 &= \sum_{m_d=1}^{t-1} (t - m_d - 1)^{d-1} \\
 &= \sum_{k=0}^{t-2} k^{d-1} = \frac{1}{d} (B_d(t-1) - B_d)
 \end{aligned}$$

From Exercises 2.15 and 2.16

$$\begin{aligned}
 L_{\mathcal{P}}(-t) &= \frac{1}{d} (B_d(-t+2) - B_d) = (-1)^d \frac{1}{d} (B_d(t-1) - B_d) \\
 &= (-1)^d L_{\mathcal{P}^\circ}(t)
 \end{aligned}$$

## Pyramids over polytopes — generalization

$\mathcal{Q}$  a convex  $(d-1)$ -polytope,  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$  the vertices of  $\mathcal{Q}$

### Definition (Pyramid over a polytope)

The **pyramid** over  $\mathcal{Q}$  is the convex hull of  $(\mathbf{v}_1, 0), (\mathbf{v}_2, 0), \dots, (\mathbf{v}_m, 0)$ , and  $(0, \dots, 0, 1)$

Denoted by  $\text{Pyr}(\mathcal{Q})$

### Note


$$\mathcal{P} = \text{Pyr}(\square)$$

### Question

What is  $L_{\text{Pyr}(\mathcal{Q})}(t) := \#(t \text{ Pyr}(\mathcal{Q}) \cap \mathbb{Z}^d)$ ?


Derivation of  $L_{\text{Pyr}(\mathcal{Q})}(t)$ 

$$L_{\text{Pyr}(\mathcal{Q})}(t) = 1 + L_Q(1) + L_Q(2) + \cdots + L_Q(t)$$



Derivation of  $L_{\text{Pyr}(\mathcal{Q})}(t)$ 

$$L_{\text{Pyr}(\mathcal{Q})}(t) = 1 + L_{\mathcal{Q}}(1) + L_{\mathcal{Q}}(2) + \cdots + L_{\mathcal{Q}}(t) = 1 + \sum_{j=1}^t L_{\mathcal{Q}}(j)$$



# The Ehrhart series of a pyramid

Theorem 2.4

$$\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) = \frac{\text{Ehr}_{\mathcal{Q}}(z)}{1-z}$$

Proof:

$$\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) = 1 + \sum_{t \geq 1} L_{\text{Pyr}(\mathcal{Q})}(t) z^t$$

# The Ehrhart series of a pyramid

## Theorem 2.4

$$\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) = \frac{\text{Ehr}_{\mathcal{Q}}(z)}{1-z}$$

Proof:

$$\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) = 1 + \sum_{t \geq 1} L_{\text{Pyr}(\mathcal{Q})}(t) z^t = 1 + \sum_{t \geq 1} \left( 1 + \sum_{j=1}^t L_{\mathcal{Q}}(j) \right) z^t$$

# The Ehrhart series of a pyramid

## Theorem 2.4

$$\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) = \frac{\text{Ehr}_{\mathcal{Q}}(z)}{1-z}$$

Proof:

$$\begin{aligned}\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) &= 1 + \sum_{t \geq 1} L_{\text{Pyr}(\mathcal{Q})}(t) z^t = 1 + \sum_{t \geq 1} \left( 1 + \sum_{j=1}^t L_{\mathcal{Q}}(j) \right) z^t \\ &= \sum_{t \geq 0} z^t + \sum_{t \geq 1} \sum_{j=1}^t L_{\mathcal{Q}}(j) z^t\end{aligned}$$

# The Ehrhart series of a pyramid

## Theorem 2.4

$$\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) = \frac{\text{Ehr}_{\mathcal{Q}}(z)}{1-z}$$

Proof:

$$\begin{aligned}\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) &= 1 + \sum_{t \geq 1} L_{\text{Pyr}(\mathcal{Q})}(t) z^t = 1 + \sum_{t \geq 1} \left( 1 + \sum_{j=1}^t L_{\mathcal{Q}}(j) \right) z^t \\ &= \sum_{t \geq 0} z^t + \sum_{t \geq 1} \sum_{j=1}^t L_{\mathcal{Q}}(j) z^t = \frac{1}{1-z} + \sum_{j \geq 1} L_{\mathcal{Q}}(j) \sum_{t \geq j} z^t\end{aligned}$$

# The Ehrhart series of a pyramid

## Theorem 2.4

$$\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) = \frac{\text{Ehr}_{\mathcal{Q}}(z)}{1-z}$$

Proof:

$$\begin{aligned}
 \text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) &= 1 + \sum_{t \geq 1} L_{\text{Pyr}(\mathcal{Q})}(t) z^t = 1 + \sum_{t \geq 1} \left( 1 + \sum_{j=1}^t L_{\mathcal{Q}}(j) \right) z^t \\
 &= \sum_{t \geq 0} z^t + \sum_{t \geq 1} \sum_{j=1}^t L_{\mathcal{Q}}(j) z^t = \frac{1}{1-z} + \sum_{j \geq 1} L_{\mathcal{Q}}(j) \sum_{t \geq j} z^t \\
 &= \frac{1}{1-z} + \sum_{j \geq 1} L_{\mathcal{Q}}(j) \frac{z^j}{1-z}
 \end{aligned}$$

# The Ehrhart series of a pyramid

## Theorem 2.4

$$\text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) = \frac{\text{Ehr}_{\mathcal{Q}}(z)}{1-z}$$

Proof:

$$\begin{aligned}
 \text{Ehr}_{\text{Pyr}(\mathcal{Q})}(z) &= 1 + \sum_{t \geq 1} L_{\text{Pyr}(\mathcal{Q})}(t) z^t = 1 + \sum_{t \geq 1} \left( 1 + \sum_{j=1}^t L_{\mathcal{Q}}(j) \right) z^t \\
 &= \sum_{t \geq 0} z^t + \sum_{t \geq 1} \sum_{j=1}^t L_{\mathcal{Q}}(j) z^t = \frac{1}{1-z} + \sum_{j \geq 1} L_{\mathcal{Q}}(j) \sum_{t \geq j} z^t \\
 &= \frac{1}{1-z} + \sum_{j \geq 1} L_{\mathcal{Q}}(j) \frac{z^j}{1-z} = \frac{1 + \sum_{j \geq 1} L_{\mathcal{Q}}(j) z^j}{1-z} \quad \square
 \end{aligned}$$

## Summary: The pyramid

Theorem 2.5

 $\mathcal{P}$  the  $d$ -pyramid

$$\mathcal{P} = \{(x_1, x_2, \dots, x_d) \in \mathbb{R}^d : 0 \leq x_1, x_2, \dots, x_{d-1} \leq 1 - x_d \leq 1\}$$

(a) The lattice-point enumerator of  $\mathcal{P}$  is the polynomial

$$L_{\mathcal{P}}(t) = \frac{1}{d} (B_d(t+2) - B_d)$$

(b) Its evaluation at negative integers yields  $(-1)^d L_{\mathcal{P}}(-t) = L_{\mathcal{P}^\circ}(t)$ (c) The Ehrhart series of  $\mathcal{P}$  is  $\text{Ehr}_{\mathcal{P}}(z) = \frac{\sum_{k=1}^{d-1} A(d-1, k) z^{k-1}}{(1-z)^{d+1}}$ 

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes

# Cross-polytopes

## Definition (Cross-polytope)


- Vertices

$$\pm \mathbf{e}_1, \pm \mathbf{e}_2, \dots, \pm \mathbf{e}_d$$

- Hyperplane description

$$\Diamond := \{(x_1, x_2, \dots, x_d) \in \mathbb{R}^d : |x_1| + |x_2| + \cdots + |x_d| \leq 1\} \quad (14)$$

## An octahedron = a 3-dimensional cross-polytope



# The bipyramid over a polytope — generalization

$\mathcal{Q}$  a convex  $(d-1)$ -polytope containing the origin,  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$  the vertices of  $\mathcal{Q}$

## Definition (bipyramid over a polytope)

The **bipyramid** over  $\mathcal{Q}$  is the convex hull of  $(\mathbf{v}_1, 0)$ ,  
 $(\mathbf{v}_2, 0), \dots, (\mathbf{v}_m, 0)$ ,  $(0, \dots, 0, 1)$ , and  $(0, \dots, 0, -1)$

Denoted by  $\text{BiPyr}(\mathcal{Q})$

## Note

$d$ -dim cross-polytope =  $\text{BiPyr}((d-1)\text{-dim cross-polytope})$

## The lattice-point enumerator and the Ehrhart series of a bipyramid

$$L_{\text{BiPyr}(\mathcal{Q})}(t) = 2 + 2L_{\mathcal{Q}}(1) + 2L_{\mathcal{Q}}(2) + \cdots + 2L_{\mathcal{Q}}(t-1) + L_{\mathcal{Q}}(t)$$

## The lattice-point enumerator and the Ehrhart series of a bipyramid

$$\begin{aligned}L_{\text{BiPyr}(\mathcal{Q})}(t) &= 2 + 2L_{\mathcal{Q}}(1) + 2L_{\mathcal{Q}}(2) + \cdots + 2L_{\mathcal{Q}}(t-1) + L_{\mathcal{Q}}(t) \\&= 2 + 2 \sum_{j=1}^{t-1} L_{\mathcal{Q}}(j) + L_{\mathcal{Q}}(t)\end{aligned}$$

## The lattice-point enumerator and the Ehrhart series of a bipyramid

$$\begin{aligned}L_{\text{BiPyr}(\mathcal{Q})}(t) &= 2 + 2L_{\mathcal{Q}}(1) + 2L_{\mathcal{Q}}(2) + \cdots + 2L_{\mathcal{Q}}(t-1) + L_{\mathcal{Q}}(t) \\&= 2 + 2 \sum_{j=1}^{t-1} L_{\mathcal{Q}}(j) + L_{\mathcal{Q}}(t)\end{aligned}$$

## Theorem 2.6

$\text{Ehr}_{\text{BiPyr}(\mathcal{Q})}(z) = \frac{1+z}{1-z} \text{Ehr}_{\mathcal{Q}}(z)$  if  $\mathcal{Q}$  contains the origin

Proof: Exercise 2.23

## Implication to cross-polytopes

- $\diamond = 0\text{-dim cross-polytope} = \{\text{origin}\} \Rightarrow$

$$\text{Ehr}_{\diamond}(z) = \frac{1}{1-z}$$

## Implication to cross-polytopes

- $\diamond = 0\text{-dim cross-polytope} = \{\text{origin}\} \Rightarrow$

$$\text{Ehr}_{\diamond}(z) = \frac{1}{1-z}$$

- $\diamond = d\text{-dim cross-polytope} \Rightarrow$

$$\text{Ehr}_{\diamond}(z) = \frac{(1+z)^d}{(1-z)^{d+1}}$$

## Implication to cross-polytopes

- $\diamond = 0\text{-dim cross-polytope} = \{\text{origin}\} \Rightarrow$

$$\text{Ehr}_{\diamond}(z) = \frac{1}{1-z}$$

- $\diamond = d\text{-dim cross-polytope} \Rightarrow$

$$\text{Ehr}_{\diamond}(z) = \frac{(1+z)^d}{(1-z)^{d+1}}$$

Let's derive  $L_{\diamond}(t)$  from  $\text{Ehr}_{\diamond}(z)!$

# Lattice-point enumerator of a cross-polytope

$$\text{Ehr}_{\diamond}(z) = \frac{(1+z)^d}{(1-z)^{d+1}}$$

# Lattice-point enumerator of a cross-polytope

$$\text{Ehr}_{\diamond}(z) = \frac{(1+z)^d}{(1-z)^{d+1}} = \frac{\sum_{k=0}^d \binom{d}{k} z^k}{(1-z)^{d+1}}$$

# Lattice-point enumerator of a cross-polytope

$$\text{Ehr}_{\diamond}(z) = \frac{(1+z)^d}{(1-z)^{d+1}} = \frac{\sum_{k=0}^d \binom{d}{k} z^k}{(1-z)^{d+1}} = \sum_{k=0}^d \binom{d}{k} z^k \sum_{t \geq 0} \binom{t+d}{d} z^t$$

## Lattice-point enumerator of a cross-polytope

$$\begin{aligned}\text{Ehr}_{\diamond}(z) &= \frac{(1+z)^d}{(1-z)^{d+1}} = \frac{\sum_{k=0}^d \binom{d}{k} z^k}{(1-z)^{d+1}} = \sum_{k=0}^d \binom{d}{k} z^k \sum_{t \geq 0} \binom{t+d}{d} z^t \\ &= \sum_{k=0}^d \binom{d}{k} \sum_{t \geq k} \binom{t-k+d}{d} z^t\end{aligned}$$

## Lattice-point enumerator of a cross-polytope

$$\begin{aligned}\text{Ehr}_{\diamond}(z) &= \frac{(1+z)^d}{(1-z)^{d+1}} = \frac{\sum_{k=0}^d \binom{d}{k} z^k}{(1-z)^{d+1}} = \sum_{k=0}^d \binom{d}{k} z^k \sum_{t \geq 0} \binom{t+d}{d} z^t \\ &= \sum_{k=0}^d \binom{d}{k} \sum_{t \geq k} \binom{t-k+d}{d} z^t \\ &= \sum_{k=0}^d \binom{d}{k} \sum_{t \geq 0} \binom{t-k+d}{d} z^t\end{aligned}$$

# Lattice-point enumerator of a cross-polytope

$$\begin{aligned}
 \text{Ehr}_{\diamond}(z) &= \frac{(1+z)^d}{(1-z)^{d+1}} = \frac{\sum_{k=0}^d \binom{d}{k} z^k}{(1-z)^{d+1}} = \sum_{k=0}^d \binom{d}{k} z^k \sum_{t \geq 0} \binom{t+d}{d} z^t \\
 &= \sum_{k=0}^d \binom{d}{k} \sum_{t \geq k} \binom{t-k+d}{d} z^t \\
 &= \sum_{k=0}^d \binom{d}{k} \sum_{t \geq 0} \binom{t-k+d}{d} z^t \\
 &= \sum_{t \geq 0} \sum_{k=0}^d \binom{d}{k} \binom{t-k+d}{d} z^t
 \end{aligned}$$

# Lattice-point enumerator of a cross-polytope

$$\begin{aligned}
 \text{Ehr}_{\diamond}(z) &= \frac{(1+z)^d}{(1-z)^{d+1}} = \frac{\sum_{k=0}^d \binom{d}{k} z^k}{(1-z)^{d+1}} = \sum_{k=0}^d \binom{d}{k} z^k \sum_{t \geq 0} \binom{t+d}{d} z^t \\
 &= \sum_{k=0}^d \binom{d}{k} \sum_{t \geq k} \binom{t-k+d}{d} z^t \\
 &= \sum_{k=0}^d \binom{d}{k} \sum_{t \geq 0} \binom{t-k+d}{d} z^t \\
 &= \sum_{t \geq 0} \sum_{k=0}^d \binom{d}{k} \binom{t-k+d}{d} z^t
 \end{aligned}$$

Therefore, for all  $t \geq 1$

$$L_{\diamond}(t) = \sum_{k=0}^d \binom{d}{k} \binom{t-k+d}{d}$$

Counting the lattice points in  $\diamond$ 

$$L_{\diamond}^{\circ}(t) = \# \{ (m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \cdots + |m_d| < t \}$$

Counting the lattice points in  $\diamond$ 

$$\begin{aligned}L_{\diamond}(t) &= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \dots + |m_d| < t\} \\&= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \dots + |m_d| \leq t-1\}\end{aligned}$$

Counting the lattice points in  $\diamondsuit^\circ$ 

$$\begin{aligned}L_{\diamondsuit^\circ}(t) &= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \dots + |m_d| < t\} \\&= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \dots + |m_d| \leq t-1\} \\&= L_{\diamondsuit}(t-1)\end{aligned}$$

Counting the lattice points in  $\diamondsuit^\circ$ 

$$\begin{aligned}L_{\diamondsuit^\circ}(t) &= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \cdots + |m_d| < t\} \\&= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \cdots + |m_d| \leq t-1\} \\&= L_{\diamondsuit}(t-1) \\&= \sum_{k=0}^d \binom{d}{d-k} \binom{t-1+d-k}{d}\end{aligned}$$

Counting the lattice points in  $\diamondsuit^\circ$ 

$$\begin{aligned}L_{\diamondsuit^\circ}(t) &= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \cdots + |m_d| < t\} \\&= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \cdots + |m_d| \leq t-1\} \\&= L_{\diamondsuit}(t-1) \\&= \sum_{k=0}^d \binom{d}{d-k} \binom{t-1+d-k}{d} \\&= (-1)^d \sum_{k=0}^d \binom{d}{k} (-1)^d \binom{t-1+k}{d}\end{aligned}$$

Counting the lattice points in  $\diamondsuit^\circ$ 

$$\begin{aligned}
 L_{\diamondsuit^\circ}(t) &= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \cdots + |m_d| < t\} \\
 &= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \cdots + |m_d| \leq t-1\} \\
 &= L_{\diamondsuit}(t-1) \\
 &= \sum_{k=0}^d \binom{d}{d-k} \binom{t-1+d-k}{d} \\
 &= (-1)^d \sum_{k=0}^d \binom{d}{k} (-1)^d \binom{t-1+k}{d} \\
 &= (-1)^d \sum_{k=0}^d \binom{d}{k} \binom{-t-k+d}{d} \quad (\text{by Ex. 2.10})
 \end{aligned}$$

Counting the lattice points in  $\diamondsuit^\circ$ 

$$\begin{aligned}
 L_{\diamondsuit^\circ}(t) &= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \cdots + |m_d| < t\} \\
 &= \#\{(m_1, m_2, \dots, m_d) \in \mathbb{Z}^d : |m_1| + |m_2| + \cdots + |m_d| \leq t-1\} \\
 &= L_{\diamondsuit}(t-1) \\
 &= \sum_{k=0}^d \binom{d}{d-k} \binom{t-1+d-k}{d} \\
 &= (-1)^d \sum_{k=0}^d \binom{d}{k} (-1)^d \binom{t-1+k}{d} \\
 &= (-1)^d \sum_{k=0}^d \binom{d}{k} \binom{-t-k+d}{d} \quad (\text{by Ex. 2.10}) \\
 &= (-1)^d L_{\diamondsuit}(-t)
 \end{aligned}$$

## Summary: Cross-polytopes

## Theorem 2.7

$\diamond$  the cross-polytope in  $\mathbb{R}^d$

(a) The lattice-point enumerator of  $\diamond$  is the polynomial

$$L_{\diamond}(t) = \sum_{k=0}^d \binom{d}{k} \binom{t-k+d}{d}$$

(b) Its evaluation at negative integers yields  $(-1)^d L_{\diamond}(-t) = L_{\diamond^\circ}(t)$

(c) The Ehrhart series of  $\mathcal{P}$  is  $\text{Ehr}_{\diamond}(z) = \frac{(1+z)^d}{(1-z)^{d+1}}$

□

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes

# Let's get back to $\mathbb{R}^2$ : Pick's theorem

## Theme

A strange connection between the number of lattice points and the area of an integral convex **polygon**


## Theorem 2.8 (Pick's theorem)

For an integral convex polygon  $\mathcal{P}$

$$A = I + \frac{1}{2}B - 1,$$

where

- $A$  = the area of  $\mathcal{P}$
- $I$  = # of lattice points in  $\mathcal{P}$
- $B$  = # of lattice points on  $\partial\mathcal{P}$



## Proof of Pick's theorem (sketch)

- $\mathcal{P}$  is partitioned into  $\mathcal{P}_1$  and  $\mathcal{P}_2 \Rightarrow$

$$I + \frac{1}{2}B - 1 = (I_1 + \frac{1}{2}B_1 - 1) + (I_2 + \frac{1}{2}B_2 - 1)$$

## Proof of Pick's theorem (sketch)

- $\mathcal{P}$  is partitioned into  $\mathcal{P}_1$  and  $\mathcal{P}_2 \Rightarrow$

$$I + \frac{1}{2}B - 1 = (I_1 + \frac{1}{2}B_1 - 1) + (I_2 + \frac{1}{2}B_2 - 1)$$



- $\therefore$  Enough to prove for triangles

## Proof of Pick's theorem (sketch)

- $\mathcal{P}$  is partitioned into  $\mathcal{P}_1$  and  $\mathcal{P}_2 \Rightarrow$

$$I + \frac{1}{2}B - 1 = (I_1 + \frac{1}{2}B_1 - 1) + (I_2 + \frac{1}{2}B_2 - 1)$$

- $\therefore$  Enough to prove for triangles
- Embed a triangle into a rectangle





## Proof of Pick's theorem (sketch)

- $\mathcal{P}$  is partitioned into  $\mathcal{P}_1$  and  $\mathcal{P}_2 \Rightarrow$

$$I + \frac{1}{2}B - 1 = (I_1 + \frac{1}{2}B_1 - 1) + (I_2 + \frac{1}{2}B_2 - 1)$$

- $\therefore$  Enough to prove for triangles
- Embed a triangle into a rectangle
- $\therefore$  Enough to prove for right triangles and rectangles





## Proof of Pick's theorem (sketch)

- $\mathcal{P}$  is partitioned into  $\mathcal{P}_1$  and  $\mathcal{P}_2 \Rightarrow$

$$I + \frac{1}{2}B - 1 = (I_1 + \frac{1}{2}B_1 - 1) + (I_2 + \frac{1}{2}B_2 - 1)$$

- $\therefore$  Enough to prove for triangles
- Embed a triangle into a rectangle
- $\therefore$  Enough to prove for right triangles and rectangles
- Ex. 2.24 will finish

□



## Summary (before the proof): an integral convex polygon

- $\#(\mathcal{P} \cap \mathbb{Z}^2) = I + B = (A - \frac{1}{2}B + 1) + B = A + \frac{1}{2}B + 1$

### Theorem 2.9

- (a) The lattice-point enumerator of  $\mathcal{P}$  is the polynomial

$$L_{\mathcal{P}}(t) = A t^2 + \frac{1}{2}B t + 1$$

- (b) Its evaluation at negative integers yields the relation

$$L_{\mathcal{P}}(-t) = L_{\mathcal{P}^\circ}(t)$$

- (c) The Ehrhart series of  $\mathcal{P}$  is

$$\text{Ehr}_{\mathcal{P}}(z) = \frac{\left(A - \frac{B}{2} + 1\right) z^2 + \left(A + \frac{B}{2} - 2\right) z + 1}{(1 - z)^3}$$

## Lattice-point enumerator of an integral convex polygon

Proof of Thm 2.9(a):

- Inflating by factor of  $t$  makes
  - the area larger by factor of  $t^2$  (Ex. 2.25)
  - the perimeter larger by factor of  $t$  (Ex. 2.25)
- Then, Pick's theorem proves □

## Lattice-point enumerator of an integral convex polygon

Proof of Thm 2.9(a):

- Inflating by factor of  $t$  makes
  - the area larger by factor of  $t^2$  (Ex. 2.25)
  - the perimeter larger by factor of  $t$  (Ex. 2.25)
- Then, Pick's theorem proves □

Proof of Thm 2.9(b):

$$L_{\mathcal{P}^\circ}(t) = L_{\mathcal{P}}(t) - B t$$

## Lattice-point enumerator of an integral convex polygon

Proof of Thm 2.9(a):

- Inflating by factor of  $t$  makes
  - the area larger by factor of  $t^2$  (Ex. 2.25)
  - the perimeter larger by factor of  $t$  (Ex. 2.25)
- Then, Pick's theorem proves □

Proof of Thm 2.9(b):

$$\begin{aligned} L_{\mathcal{P}^\circ}(t) &= L_{\mathcal{P}}(t) - B t \\ &= \left( A t^2 + \frac{1}{2} B t + 1 \right) - B t \end{aligned}$$

## Lattice-point enumerator of an integral convex polygon

Proof of Thm 2.9(a):

- Inflating by factor of  $t$  makes
  - the area larger by factor of  $t^2$  (Ex. 2.25)
  - the perimeter larger by factor of  $t$  (Ex. 2.25)
- Then, Pick's theorem proves □

Proof of Thm 2.9(b):

$$\begin{aligned}
 L_{\mathcal{P}^\circ}(t) &= L_{\mathcal{P}}(t) - B t \\
 &= \left( A t^2 + \frac{1}{2} B t + 1 \right) - B t \\
 &= A t^2 - \frac{1}{2} B t + 1 = L_{\mathcal{P}}(-t) \quad \square
 \end{aligned}$$

## Ehrhart series of an integral convex polygon

Proof of Thm 2.9(c):

$$\text{Ehr}_{\mathcal{P}}(z) = 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t$$

## Ehrhart series of an integral convex polygon

Proof of Thm 2.9(c):

$$\begin{aligned}\text{Ehr}_{\mathcal{P}}(z) &= 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t \\ &= \sum_{t \geq 0} \left( A t^2 + \frac{B}{2} t + 1 \right) z^t\end{aligned}$$

## Ehrhart series of an integral convex polygon

Proof of Thm 2.9(c):

$$\begin{aligned}\text{Ehr}_{\mathcal{P}}(z) &= 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t \\ &= \sum_{t \geq 0} \left( A t^2 + \frac{B}{2} t + 1 \right) z^t \\ &= A \frac{z^2 + z}{(1 - z)^3} + \frac{B}{2} \frac{z}{(1 - z)^2} + \frac{1}{1 - z}\end{aligned}$$

## Ehrhart series of an integral convex polygon


Proof of Thm 2.9(c):

$$\begin{aligned}
 \text{Ehr}_{\mathcal{P}}(z) &= 1 + \sum_{t \geq 1} L_{\mathcal{P}}(t) z^t \\
 &= \sum_{t \geq 0} \left( A t^2 + \frac{B}{2} t + 1 \right) z^t \\
 &= A \frac{z^2 + z}{(1-z)^3} + \frac{B}{2} \frac{z}{(1-z)^2} + \frac{1}{1-z} \\
 &= \frac{\left( A - \frac{B}{2} + 1 \right) z^2 + \left( A + \frac{B}{2} - 2 \right) z + 1}{(1-z)^3} \quad \square
 \end{aligned}$$

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes


# Roadmap

## Contents



# Roadmap

## Contents



## Goal of this section

- Develop a theory for rational convex polygons
- Introduce a quasipolynomial

# Steps towards rational convex polygons

- Triangulate a rational convex polygon
- → Enough to study triangles


# Steps towards rational convex polygons

- Triangulate a rational convex polygon
- → Enough to study triangles
- Embed a triangle into a rectangle
- → Enough to study right triangles

# Steps towards rational convex polygons

- Triangulate a rational convex polygon
- → Enough to study triangles
- Embed a triangle into a rectangle
- → Enough to study right triangles
- Translate, rotate, and mirror a right triangle
- → Enough to study the following type of triangles

## A right triangle: setup



$$\mathcal{T} = \left\{ (x, y) \in \mathbb{R}^2 : x \geq \frac{a}{d}, y \geq \frac{b}{d}, ex + fy \leq r \right\} \quad (16)$$

- $a, b, d, e, f, r \in \mathbb{Z}_{\geq 0}$ ,  $ea + fb \leq rd$ ,  $a, b < d$
- For brevity,  $e, f$  coprime

## Lattice-point enumerator: introducing a slack variable

$$L_T(t) = \# \left\{ (m, n) \in \mathbb{Z}^2 : m \geq \frac{ta}{d}, n \geq \frac{tb}{d}, em + fn \leq tr \right\}$$

## Lattice-point enumerator: introducing a slack variable

$$\begin{aligned} L_T(t) &= \# \left\{ (m, n) \in \mathbb{Z}^2 : m \geq \frac{ta}{d}, n \geq \frac{tb}{d}, em + fn \leq tr \right\} \\ &= \# \left\{ (m, n, s) \in \mathbb{Z}^3 : m \geq \frac{ta}{d}, n \geq \frac{tb}{d}, s \geq 0, em + fn + s = tr \right\} \end{aligned}$$

## Lattice-point enumerator: introducing a slack variable

$$\begin{aligned} L_T(t) &= \# \left\{ (m, n) \in \mathbb{Z}^2 : m \geq \frac{ta}{d}, n \geq \frac{tb}{d}, em + fn \leq tr \right\} \\ &= \# \left\{ (m, n, s) \in \mathbb{Z}^3 : m \geq \frac{ta}{d}, n \geq \frac{tb}{d}, s \geq 0, em + fn + s = tr \right\} \end{aligned}$$

This is interpreted as the coefficient of  $z^{tr}$  in the function

$$\left( \sum_{m \geq \frac{ta}{d}} z^{em} \right) \left( \sum_{n \geq \frac{tb}{d}} z^{fn} \right) \left( \sum_{s \geq 0} z^s \right),$$

where the subscript under a summation sign means “sum over all integers satisfying this condition”

## Lattice-point enumerator: a power series

$$\left( \sum_{m \geq \lceil \frac{ta}{d} \rceil} z^{em} \right) \left( \sum_{n \geq \lceil \frac{tb}{d} \rceil} z^{fn} \right) \left( \sum_{s \geq 0} z^s \right)$$

(17)

## Lattice-point enumerator: a power series

$$\left( \sum_{m \geq \lceil \frac{ta}{d} \rceil} z^{em} \right) \left( \sum_{n \geq \lceil \frac{tb}{d} \rceil} z^{fn} \right) \left( \sum_{s \geq 0} z^s \right) = \frac{z^{\lceil \frac{ta}{d} \rceil e}}{1 - z^e} \frac{z^{\lceil \frac{tb}{d} \rceil f}}{1 - z^f} \frac{1}{1 - z}$$

(17)

## Lattice-point enumerator: a power series

$$\begin{aligned}
 & \left( \sum_{m \geq \lceil \frac{ta}{d} \rceil} z^{em} \right) \left( \sum_{n \geq \lceil \frac{tb}{d} \rceil} z^{fn} \right) \left( \sum_{s \geq 0} z^s \right) = \frac{z^{\lceil \frac{ta}{d} \rceil e}}{1 - z^e} \frac{z^{\lceil \frac{tb}{d} \rceil f}}{1 - z^f} \frac{1}{1 - z} \\
 &= \frac{z^{u+v}}{(1 - z^e)(1 - z^f)(1 - z)}, \tag{17}
 \end{aligned}$$

where

$$u := \left\lceil \frac{ta}{d} \right\rceil e \quad \text{and} \quad v := \left\lceil \frac{tb}{d} \right\rceil f \tag{18}$$

## Lattice-point enumerator: a power series

$$\begin{aligned}
 & \left( \sum_{m \geq \lceil \frac{ta}{d} \rceil} z^{em} \right) \left( \sum_{n \geq \lceil \frac{tb}{d} \rceil} z^{fn} \right) \left( \sum_{s \geq 0} z^s \right) = \frac{z^{\lceil \frac{ta}{d} \rceil e}}{1 - z^e} \frac{z^{\lceil \frac{tb}{d} \rceil f}}{1 - z^f} \frac{1}{1 - z} \\
 & = \frac{z^{u+v}}{(1 - z^e)(1 - z^f)(1 - z)}, \tag{17}
 \end{aligned}$$

where

$$u := \left\lceil \frac{ta}{d} \right\rceil e \quad \text{and} \quad v := \left\lceil \frac{tb}{d} \right\rceil f \tag{18}$$

Therefore,

$$L_T(t) = \text{const} \left( \frac{z^{u+v-tr}}{(1 - z^e)(1 - z^f)(1 - z)} \right)$$

## Lattice-point enumerator: a power series

$$\begin{aligned}
 & \left( \sum_{m \geq \lceil \frac{ta}{d} \rceil} z^{em} \right) \left( \sum_{n \geq \lceil \frac{tb}{d} \rceil} z^{fn} \right) \left( \sum_{s \geq 0} z^s \right) = \frac{z^{\lceil \frac{ta}{d} \rceil e}}{1 - z^e} \frac{z^{\lceil \frac{tb}{d} \rceil f}}{1 - z^f} \frac{1}{1 - z} \\
 & = \frac{z^{u+v}}{(1 - z^e)(1 - z^f)(1 - z)}, \tag{17}
 \end{aligned}$$

where

$$u := \left\lceil \frac{ta}{d} \right\rceil e \quad \text{and} \quad v := \left\lceil \frac{tb}{d} \right\rceil f \tag{18}$$

Therefore,

$$\begin{aligned}
 L_T(t) &= \text{const} \left( \frac{z^{u+v-tr}}{(1 - z^e)(1 - z^f)(1 - z)} \right) \\
 &= \text{const} \left( \frac{1}{(1 - z^e)(1 - z^f)(1 - z)z^{tr-u-v}} \right)
 \end{aligned}$$

## Lattice-point enumerator: theorem

$$L_T(t) = \text{const} \left( \frac{1}{(1-z^e)(1-z^f)(1-z)z^{tr-u-v}} \right)$$

## Lattice-point enumerator: theorem

$$L_T(t) = \text{const} \left( \frac{1}{(1-z^e)(1-z^f)(1-z)z^{tr-u-v}} \right)$$

- Note:  $u + v - tr - e - f - 1 < 0$  (Ex. 2.31)

## Lattice-point enumerator: theorem

$$L_{\mathcal{T}}(t) = \text{const} \left( \frac{1}{(1-z^e)(1-z^f)(1-z)z^{tr-u-v}} \right)$$

- Note:  $u + v - tr - e - f - 1 < 0$  (Ex. 2.31)
- A calculation gives the following theorem (Ex. 2.32)

## Theorem 2.10

For the triangle  $\mathcal{T}$  given by (16), where  $e$  and  $f$  are coprime,

$$\begin{aligned} L_{\mathcal{T}}(t) &= \frac{1}{2ef} (tr - u - v)^2 + \frac{1}{2} (tr - u - v) \left( \frac{1}{e} + \frac{1}{f} + \frac{1}{ef} \right) \\ &\quad + \frac{1}{4} \left( 1 + \frac{1}{e} + \frac{1}{f} \right) + \frac{1}{12} \left( \frac{e}{f} + \frac{f}{e} + \frac{1}{ef} \right) \\ &\quad + \frac{1}{e} \sum_{j=1}^{e-1} \frac{\xi_e^{j(v-tr)}}{(1 - \xi_e^{jf})(1 - \xi_e^j)} + \frac{1}{f} \sum_{l=1}^{f-1} \frac{\xi_f^{l(u-tr)}}{(1 - \xi_f^{le})(1 - \xi_f^l)} \end{aligned} \quad \square$$

Properties of this  $L_T(t)$ 

$$\begin{aligned}
 L_T(t) = & \frac{1}{2ef} (tr - u - v)^2 + \frac{1}{2} (tr - u - v) \left( \frac{1}{e} + \frac{1}{f} + \frac{1}{ef} \right) \\
 & + \frac{1}{4} \left( 1 + \frac{1}{e} + \frac{1}{f} \right) + \frac{1}{12} \left( \frac{e}{f} + \frac{f}{e} + \frac{1}{ef} \right) \\
 & + \frac{1}{e} \sum_{j=1}^{e-1} \frac{\xi_e^{j(v-tr)}}{(1 - \xi_e^{jf})(1 - \xi_e^j)} + \frac{1}{f} \sum_{l=1}^{f-1} \frac{\xi_f^{l(u-tr)}}{(1 - \xi_f^{le})(1 - \xi_f^l)}
 \end{aligned}$$

Properties of this  $L_T(t)$ 

$$\begin{aligned}
 L_T(t) = & \frac{1}{2ef} (tr - u - v)^2 + \frac{1}{2} (tr - u - v) \left( \frac{1}{e} + \frac{1}{f} + \frac{1}{ef} \right) \\
 & + \frac{1}{4} \left( 1 + \frac{1}{e} + \frac{1}{f} \right) + \frac{1}{12} \left( \frac{e}{f} + \frac{f}{e} + \frac{1}{ef} \right) \\
 & + \frac{1}{e} \sum_{j=1}^{e-1} \frac{\xi_e^{j(v-tr)}}{(1 - \xi_e^{jf})(1 - \xi_e^j)} + \frac{1}{f} \sum_{l=1}^{f-1} \frac{\xi_f^{l(u-tr)}}{(1 - \xi_f^{le})(1 - \xi_f^l)}
 \end{aligned}$$

- $L_T(t)$  is a **quadratic** fn if we forget the last two sums and  $u, v$

Properties of this  $L_T(t)$ 

$$\begin{aligned}
 L_T(t) = & \frac{1}{2ef} (tr - u - v)^2 + \frac{1}{2} (tr - u - v) \left( \frac{1}{e} + \frac{1}{f} + \frac{1}{ef} \right) \\
 & + \frac{1}{4} \left( 1 + \frac{1}{e} + \frac{1}{f} \right) + \frac{1}{12} \left( \frac{e}{f} + \frac{f}{e} + \frac{1}{ef} \right) \\
 & + \frac{1}{e} \sum_{j=1}^{e-1} \frac{\xi_e^{j(v-tr)}}{(1 - \xi_e^{jf})(1 - \xi_e^j)} + \frac{1}{f} \sum_{l=1}^{f-1} \frac{\xi_f^{l(u-tr)}}{(1 - \xi_f^{le})(1 - \xi_f^l)}
 \end{aligned}$$

- $L_T(t)$  is a **quadratic** fn if we forget the last two sums and  $u, v$
- the last two sums are **periodic**

Properties of this  $L_T(t)$ 

$$\begin{aligned}
 L_T(t) = & \frac{1}{2ef} (tr - u - v)^2 + \frac{1}{2} (tr - u - v) \left( \frac{1}{e} + \frac{1}{f} + \frac{1}{ef} \right) \\
 & + \frac{1}{4} \left( 1 + \frac{1}{e} + \frac{1}{f} \right) + \frac{1}{12} \left( \frac{e}{f} + \frac{f}{e} + \frac{1}{ef} \right) \\
 & + \frac{1}{e} \sum_{j=1}^{e-1} \frac{\xi_e^{j(v-tr)}}{(1 - \xi_e^{jf})(1 - \xi_e^j)} + \frac{1}{f} \sum_{l=1}^{f-1} \frac{\xi_f^{l(u-tr)}}{(1 - \xi_f^{le})(1 - \xi_f^l)}
 \end{aligned}$$

- $L_T(t)$  is a **quadratic** fn if we forget the last two sums and  $u, v$
- the last two sums are **periodic**
- $u = \lceil \frac{ta}{d} \rceil e$  and  $v = \lceil \frac{tb}{d} \rceil f$  show **periodic** behaviors

Properties of this  $L_T(t)$ 

$$\begin{aligned}
 L_T(t) = & \frac{1}{2ef} (tr - u - v)^2 + \frac{1}{2} (tr - u - v) \left( \frac{1}{e} + \frac{1}{f} + \frac{1}{ef} \right) \\
 & + \frac{1}{4} \left( 1 + \frac{1}{e} + \frac{1}{f} \right) + \frac{1}{12} \left( \frac{e}{f} + \frac{f}{e} + \frac{1}{ef} \right) \\
 & + \frac{1}{e} \sum_{j=1}^{e-1} \frac{\xi_e^{j(v-tr)}}{(1 - \xi_e^{jf})(1 - \xi_e^j)} + \frac{1}{f} \sum_{l=1}^{f-1} \frac{\xi_f^{l(u-tr)}}{(1 - \xi_f^{le})(1 - \xi_f^l)}
 \end{aligned}$$

- $L_T(t)$  is a **quadratic** fn if we forget the last two sums and  $u, v$
- the last two sums are **periodic**
- $u = \lceil \frac{ta}{d} \rceil e$  and  $v = \lceil \frac{tb}{d} \rceil f$  show **periodic** behaviors

Therefore,  $L_T(t)$  is a “quadratic polynomial” in  $t$  whose coefficients are periodic in  $t$

# Quasipolynomials

## Definition (Quasipolynomial)

A function  $Q$  in  $t$  is **quasipolynomial** if  $Q$  can be expressed as

$$Q(t) = c_n(t) t^n + \cdots + c_1(t) t + c_0(t),$$

where  $c_0, \dots, c_n$  are periodic functions in  $t$

- The **degree** of  $Q$  is  $n$  (assuming that  $c_n$  is not the zero function)
- The **period** of  $Q$  is the least common period of  $c_0, \dots, c_n$

## Constituents of a quasipolynomial

$Q$  a quasipolynomial in  $t$

- $\exists k$  and polynomials  $p_0, p_1, \dots, p_{k-1}$  s.t.

$$Q(t) = \begin{cases} p_0(t) & \text{if } t \equiv 0 \pmod{k}, \\ p_1(t) & \text{if } t \equiv 1 \pmod{k}, \\ \vdots & \\ p_{k-1}(t) & \text{if } t \equiv k-1 \pmod{k} \end{cases}$$

- The minimal such  $k$  is the period of  $Q$

## Definition (Constituent)

For this minimal  $k$ , the polynomials  $p_0, p_1, \dots, p_{k-1}$  are the **constituents** of  $Q$

## The lattice-point enumerator of a rational polygon is a quasipolynomial

$$\begin{aligned}
 L_T(t) = & \frac{1}{2ef} (tr - u - v)^2 + \frac{1}{2} (tr - u - v) \left( \frac{1}{e} + \frac{1}{f} + \frac{1}{ef} \right) \\
 & + \frac{1}{4} \left( 1 + \frac{1}{e} + \frac{1}{f} \right) + \frac{1}{12} \left( \frac{e}{f} + \frac{f}{e} + \frac{1}{ef} \right) \\
 & + \frac{1}{e} \sum_{j=1}^{e-1} \frac{\xi_e^{j(v-tr)}}{(1 - \xi_e^{jf})(1 - \xi_e^j)} + \frac{1}{f} \sum_{l=1}^{f-1} \frac{\xi_f^{l(u-tr)}}{(1 - \xi_f^{le})(1 - \xi_f^l)}
 \end{aligned}$$

- This is a quasipolynomial of degree 2

## Theorem 2.11


$\mathcal{P}$  any rational polygon  $\Rightarrow$

- $L_{\mathcal{P}}(t)$  is a quasipolynomial of degree 2
- Its leading coefficient is the area of  $\mathcal{P}$  (in particular, a constant)

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes

# Roadmap

## Contents



## Goal of this section

- Develop a theory for rational convex polytopes

## Rational polytopes: setup

### Definition (recap)

A polytope  $\mathcal{P}$  is **rational** if all of its vertices have rational coordinates

We are interested in  $\#(t \mathcal{P} \cap \mathbb{Z}^d)$

## Rational polytopes: setup

### Definition (recap)

A polytope  $\mathcal{P}$  is **rational** if all of its vertices have rational coordinates

We are interested in  $\#(t\mathcal{P} \cap \mathbb{Z}^d)$

- Consider a hyperplane description of  $\mathcal{P}$

## Rational polytopes: setup

### Definition (recap)

A polytope  $\mathcal{P}$  is **rational** if all of its vertices have rational coordinates

We are interested in  $\#(t \mathcal{P} \cap \mathbb{Z}^d)$

- Consider a hyperplane description of  $\mathcal{P}$
- Every coefficient can be chosen as an integer (Ex. 2.7)

## Rational polytopes: setup

### Definition (recap)

A polytope  $\mathcal{P}$  is **rational** if all of its vertices have rational coordinates

We are interested in  $\#(t\mathcal{P} \cap \mathbb{Z}^d)$

- Consider a hyperplane description of  $\mathcal{P}$
- Every coefficient can be chosen as an integer (Ex. 2.7)
- Inequalities are transformed into equalities (by slack var's)

## Rational polytopes: setup

### Definition (recap)

A polytope  $\mathcal{P}$  is **rational** if all of its vertices have rational coordinates

We are interested in  $\#(t\mathcal{P} \cap \mathbb{Z}^d)$

- Consider a hyperplane description of  $\mathcal{P}$
- Every coefficient can be chosen as an integer (Ex. 2.7)
- Inequalities are transformed into equalities (by slack var's)
- All pts in  $\mathcal{P}$  have nonnegative coord's (by translation)

## Rational polytopes: setup

## Definition (recap)

A polytope  $\mathcal{P}$  is **rational** if all of its vertices have rational coordinates

We are interested in  $\#(t \mathcal{P} \cap \mathbb{Z}^d)$

- Consider a hyperplane description of  $\mathcal{P}$
- Every coefficient can be chosen as an integer (Ex. 2.7)
- Inequalities are transformed into equalities (by slack var's)
- All pts in  $\mathcal{P}$  have nonnegative coord's (by translation)

Therefore, any rational polytope  $\mathcal{P}$  is expressed as

$$\mathcal{P} = \{\mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b}\} \quad (23)$$

for some integral matrix  $\mathbf{A} \in \mathbb{Z}^{m \times d}$  and some integer vector  $\mathbf{b} \in \mathbb{Z}^m$

Note:  $d$  is not necessarily the dimension of  $\mathcal{P}$

## Lattice-point enumerator of a rational polytope

$$\mathcal{P} = \{\mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b}\}$$

## Lattice-point enumerator of a rational polytope

$$\mathcal{P} = \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b} \right\}$$

Therefore,

$$t\mathcal{P} = \left\{ t\mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b} \right\}$$

## Lattice-point enumerator of a rational polytope

$$\mathcal{P} = \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b} \right\}$$

Therefore,

$$\begin{aligned} t\mathcal{P} &= \left\{ t\mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b} \right\} \\ &= \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A} \frac{\mathbf{x}}{t} = \mathbf{b} \right\} \end{aligned}$$

## Lattice-point enumerator of a rational polytope

$$\mathcal{P} = \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b} \right\}$$

Therefore,

$$\begin{aligned} t\mathcal{P} &= \left\{ t\mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b} \right\} \\ &= \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A} \frac{\mathbf{x}}{t} = \mathbf{b} \right\} \\ &= \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = t\mathbf{b} \right\} \end{aligned}$$

## Lattice-point enumerator of a rational polytope

$$\mathcal{P} = \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b} \right\}$$


Therefore,

$$\begin{aligned} t\mathcal{P} &= \left\{ t\mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = \mathbf{b} \right\} \\ &= \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A} \frac{\mathbf{x}}{t} = \mathbf{b} \right\} \\ &= \left\{ \mathbf{x} \in \mathbb{R}_{\geq 0}^d : \mathbf{A}\mathbf{x} = t\mathbf{b} \right\} \end{aligned}$$

Namely,

$$L_{\mathcal{P}}(t) = \# \left\{ \mathbf{x} \in \mathbb{Z}_{\geq 0}^d : \mathbf{A}\mathbf{x} = t\mathbf{b} \right\} \quad (24)$$

## Example: Setup



$$\mathcal{P} = \left\{ (x_1, x_2) \in \mathbb{R}^2 : x_1, x_2 \geq 0, \begin{array}{l} x_1 + 2x_2 \leq 3, \\ x_1 + x_2 \leq 2 \end{array} \right\}$$

## Example: Lattice-point enumerator

$$L_{\mathcal{P}}(t) = \# \left\{ (x_1, x_2) \in \mathbb{Z}^2 : x_1, x_2 \geq 0, \begin{array}{l} x_1 + 2x_2 \leq 3t, \\ x_1 + x_2 \leq 2t \end{array} \right\}$$

## Example: Lattice-point enumerator

$$\begin{aligned}L_{\mathcal{P}}(t) &= \# \left\{ (x_1, x_2) \in \mathbb{Z}^2 : x_1, x_2 \geq 0, \begin{array}{l} x_1 + 2x_2 \leq 3t, \\ x_1 + x_2 \leq 2t \end{array} \right\} \\&= \# \left\{ (x_1, x_2, x_3, x_4) \in \mathbb{Z}^4 : \begin{array}{l} x_1, x_2, x_3, x_4 \geq 0, \\ x_1 + 2x_2 + x_3 = 3t, \\ x_1 + x_2 + x_4 = 2t \end{array} \right\}\end{aligned}$$

## Example: Lattice-point enumerator

$$\begin{aligned}
 L_{\mathcal{P}}(t) &= \# \left\{ (x_1, x_2) \in \mathbb{Z}^2 : x_1, x_2 \geq 0, \begin{array}{l} x_1 + 2x_2 \leq 3t, \\ x_1 + x_2 \leq 2t \end{array} \right\} \\
 &= \# \left\{ (x_1, x_2, x_3, x_4) \in \mathbb{Z}^4 : \begin{array}{l} x_1, x_2, x_3, x_4 \geq 0, \\ x_1 + 2x_2 + x_3 = 3t, \\ x_1 + x_2 + x_4 = 2t \end{array} \right\} \\
 &= \# \left\{ \mathbf{x} \in \mathbb{Z}_{\geq 0}^4 : \left( \begin{array}{cccc} 1 & 2 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{array} \right) \mathbf{x} = \left( \begin{array}{c} 3t \\ 2t \end{array} \right) \right\}
 \end{aligned}$$

## Example: Power series

$$f(z_1, z_2) := \frac{1}{(1 - z_1 z_2)(1 - z_1^2 z_2)(1 - z_1)(1 - z_2) z_1^{3t} z_2^{2t}}$$

## Example: Power series

$$\begin{aligned}f(z_1, z_2) &:= \frac{1}{(1 - z_1 z_2)(1 - z_1^2 z_2)(1 - z_1)(1 - z_2)z_1^{3t} z_2^{2t}} \\&= \left( \sum_{n_1 \geq 0} (z_1 z_2)^{n_1} \right) \left( \sum_{n_2 \geq 0} (z_1^2 z_2)^{n_2} \right) \left( \sum_{n_3 \geq 0} z_1^{n_3} \right) \left( \sum_{n_4 \geq 0} z_2^{n_4} \right) \frac{1}{z_1^{3t} z_2^{2t}}\end{aligned}$$

## Example: Power series

$$\begin{aligned}
 f(z_1, z_2) &:= \frac{1}{(1 - z_1 z_2)(1 - z_1^2 z_2)(1 - z_1)(1 - z_2)z_1^{3t} z_2^{2t}} \\
 &= \left( \sum_{n_1 \geq 0} (z_1 z_2)^{n_1} \right) \left( \sum_{n_2 \geq 0} (z_1^2 z_2)^{n_2} \right) \left( \sum_{n_3 \geq 0} z_1^{n_3} \right) \left( \sum_{n_4 \geq 0} z_2^{n_4} \right) \frac{1}{z_1^{3t} z_2^{2t}} \\
 &= \sum_{n_1, \dots, n_4 \geq 0} z_1^{n_1 + 2n_2 + n_3 - 3t} z_2^{n_1 + n_2 + n_4 - 2t}
 \end{aligned}$$

## Example: Power series

$$\begin{aligned}
 f(z_1, z_2) &:= \frac{1}{(1 - z_1 z_2)(1 - z_1^2 z_2)(1 - z_1)(1 - z_2)z_1^{3t} z_2^{2t}} \\
 &= \left( \sum_{n_1 \geq 0} (z_1 z_2)^{n_1} \right) \left( \sum_{n_2 \geq 0} (z_1^2 z_2)^{n_2} \right) \left( \sum_{n_3 \geq 0} z_1^{n_3} \right) \left( \sum_{n_4 \geq 0} z_2^{n_4} \right) \frac{1}{z_1^{3t} z_2^{2t}} \\
 &= \sum_{n_1, \dots, n_4 \geq 0} z_1^{n_1 + 2n_2 + n_3 - 3t} z_2^{n_1 + n_2 + n_4 - 2t}
 \end{aligned}$$

Therefore,

$$L_{\mathcal{P}}(t) = \underset{z_1, z_2}{\text{const}} f(z_1, z_2)$$

## Example: Power series

$$\begin{aligned}
 f(z_1, z_2) &:= \frac{1}{(1 - z_1 z_2)(1 - z_1^2 z_2)(1 - z_1)(1 - z_2)z_1^{3t} z_2^{2t}} \\
 &= \left( \sum_{n_1 \geq 0} (z_1 z_2)^{n_1} \right) \left( \sum_{n_2 \geq 0} (z_1^2 z_2)^{n_2} \right) \left( \sum_{n_3 \geq 0} z_1^{n_3} \right) \left( \sum_{n_4 \geq 0} z_2^{n_4} \right) \frac{1}{z_1^{3t} z_2^{2t}} \\
 &= \sum_{n_1, \dots, n_4 \geq 0} z_1^{n_1+2n_2+n_3-3t} z_2^{n_1+n_2+n_4-2t}
 \end{aligned}$$

Therefore,

$$L_{\mathcal{P}}(t) = \underset{z_1, z_2}{\text{const}} f(z_1, z_2)$$

Then, we have (Ex. 2.36)

$$\frac{7}{4} t^2 + \frac{5}{2} t + \frac{7 + (-1)^t}{8}$$

## General case: Lattice-point enumerator and power series

## Reminder

$$L_{\mathcal{P}}(t) = \#\left\{\mathbf{x} \in \mathbb{Z}_{\geq 0}^d : \mathbf{A}\mathbf{x} = t\mathbf{b}\right\}$$

Let

- $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_d$  the columns of  $\mathbf{A}$
- $\mathbf{z} = (z_1, z_2, \dots, z_m)$

## General case: Lattice-point enumerator and power series

## Reminder

$$L_{\mathcal{P}}(t) = \#\left\{\mathbf{x} \in \mathbb{Z}_{\geq 0}^d : \mathbf{A}\mathbf{x} = t\mathbf{b}\right\}$$

Let

- $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_d$  the columns of  $\mathbf{A}$
- $\mathbf{z} = (z_1, z_2, \dots, z_m)$

Let

$$f(\mathbf{z}) = \frac{1}{(1 - \mathbf{z}^{\mathbf{c}_1})(1 - \mathbf{z}^{\mathbf{c}_2}) \cdots (1 - \mathbf{z}^{\mathbf{c}_d}) \mathbf{z}^{t\mathbf{b}}} \quad (25)$$

where  $\mathbf{z}^{\mathbf{c}} := z_1^{c_1} z_2^{c_2} \cdots z_m^{c_m}$  for  $\mathbf{c} = (c_1, c_2, \dots, c_m) \in \mathbb{Z}^m$

## General case: Lattice-point enumerator and power series

## Reminder

$$L_{\mathcal{P}}(t) = \#\left\{\mathbf{x} \in \mathbb{Z}_{\geq 0}^d : \mathbf{A}\mathbf{x} = t\mathbf{b}\right\}$$

Let

- $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_d$  the columns of  $\mathbf{A}$
- $\mathbf{z} = (z_1, z_2, \dots, z_m)$

Let

$$\begin{aligned} f(\mathbf{z}) &= \frac{1}{(1 - \mathbf{z}^{\mathbf{c}_1})(1 - \mathbf{z}^{\mathbf{c}_2}) \cdots (1 - \mathbf{z}^{\mathbf{c}_d}) \mathbf{z}^{t\mathbf{b}}} \\ &= \left( \sum_{n_1 \geq 0} \mathbf{z}^{n_1 \mathbf{c}_1} \right) \left( \sum_{n_2 \geq 0} \mathbf{z}^{n_2 \mathbf{c}_2} \right) \cdots \left( \sum_{n_d \geq 0} \mathbf{z}^{n_d \mathbf{c}_d} \right) \frac{1}{\mathbf{z}^{t\mathbf{b}}}, \end{aligned} \tag{25}$$

where  $\mathbf{z}^{\mathbf{c}} := z_1^{c_1} z_2^{c_2} \cdots z_m^{c_m}$  for  $\mathbf{c} = (c_1, c_2, \dots, c_m) \in \mathbb{Z}^m$

## General case: A typical term

$$f(\mathbf{z}) = \left( \sum_{n_1 \geq 0} \mathbf{z}^{n_1 \mathbf{c}_1} \right) \left( \sum_{n_2 \geq 0} \mathbf{z}^{n_2 \mathbf{c}_2} \right) \cdots \left( \sum_{n_d \geq 0} \mathbf{z}^{n_d \mathbf{c}_d} \right) \frac{1}{\mathbf{z}^{t\mathbf{b}}}$$

- The exponent of a typical term looks like

$$n_1 \mathbf{c}_1 + n_2 \mathbf{c}_2 + \cdots + n_d \mathbf{c}_d - t\mathbf{b} = \mathbf{A}\mathbf{n} - t\mathbf{b},$$

where  $\mathbf{n} = (n_1, n_2, \dots, n_d) \in \mathbb{Z}_{\geq 0}^d$

## General case: A typical term

$$f(\mathbf{z}) = \left( \sum_{n_1 \geq 0} \mathbf{z}^{n_1 \mathbf{c}_1} \right) \left( \sum_{n_2 \geq 0} \mathbf{z}^{n_2 \mathbf{c}_2} \right) \cdots \left( \sum_{n_d \geq 0} \mathbf{z}^{n_d \mathbf{c}_d} \right) \frac{1}{\mathbf{z}^{t\mathbf{b}}}$$

- The exponent of a typical term looks like

$$n_1 \mathbf{c}_1 + n_2 \mathbf{c}_2 + \cdots + n_d \mathbf{c}_d - t\mathbf{b} = \mathbf{An} - t\mathbf{b},$$

where  $\mathbf{n} = (n_1, n_2, \dots, n_d) \in \mathbb{Z}_{\geq 0}^d$

- Therefore, the constant term of  $f(\mathbf{z})$  counts the number of solutions  $\mathbf{n}$  to

$$\mathbf{An} - t\mathbf{b} = \mathbf{0},$$

namely, the number of lattice points in  $t\mathcal{P}$

## General case: Euler's generating function

## Theorem 2.13

Suppose the rational polytope  $\mathcal{P}$  is given by (23).

Then the lattice-point enumerator of  $\mathcal{P}$  can be computed as

$$L_{\mathcal{P}}(t) = \underset{\mathbf{z}}{\text{const}} \left( \frac{1}{(1 - \mathbf{z}^{\mathbf{c}_1})(1 - \mathbf{z}^{\mathbf{c}_2}) \cdots (1 - \mathbf{z}^{\mathbf{c}_d}) \mathbf{z}^{t\mathbf{b}}} \right) \quad \square$$

## General case: Ehrhart series

## Corollary 2.14

Suppose the rational polytope  $\mathcal{P}$  is given by (23).

Then the Ehrhart series of  $\mathcal{P}$  can be computed as

$$\text{Ehr}_{\mathcal{P}}(x) = \underset{z}{\text{const}} \left( \frac{1}{(1 - z^{c_1})(1 - z^{c_2}) \cdots (1 - z^{c_d})(1 - \frac{x}{z^b})} \right)$$

Proof:

$$\text{Ehr}_{\mathcal{P}}(x) = \sum_{t \geq 0} \underset{z}{\text{const}} \left( \frac{1}{(1 - z^{c_1})(1 - z^{c_2}) \cdots (1 - z^{c_d}) z^{t b}} \right) x^t$$

## General case: Ehrhart series

## Corollary 2.14

Suppose the rational polytope  $\mathcal{P}$  is given by (23).  
 Then the Ehrhart series of  $\mathcal{P}$  can be computed as

$$\text{Ehr}_{\mathcal{P}}(x) = \underset{z}{\text{const}} \left( \frac{1}{(1 - z^{c_1})(1 - z^{c_2}) \cdots (1 - z^{c_d})(1 - \frac{x}{z^b})} \right)$$

Proof:

$$\begin{aligned} \text{Ehr}_{\mathcal{P}}(x) &= \sum_{t \geq 0} \underset{z}{\text{const}} \left( \frac{1}{(1 - z^{c_1})(1 - z^{c_2}) \cdots (1 - z^{c_d}) z^{tb}} \right) x^t \\ &= \underset{z}{\text{const}} \left( \frac{1}{(1 - z^{c_1})(1 - z^{c_2}) \cdots (1 - z^{c_d})} \sum_{t \geq 0} \frac{x^t}{z^{tb}} \right) \end{aligned}$$

## General case: Ehrhart series

## Corollary 2.14

Suppose the rational polytope  $\mathcal{P}$  is given by (23).  
 Then the Ehrhart series of  $\mathcal{P}$  can be computed as

$$\text{Ehr}_{\mathcal{P}}(x) = \underset{z}{\text{const}} \left( \frac{1}{(1 - z^{c_1})(1 - z^{c_2}) \cdots (1 - z^{c_d})(1 - \frac{x}{z^b})} \right)$$

Proof:

$$\begin{aligned} \text{Ehr}_{\mathcal{P}}(x) &= \sum_{t \geq 0} \underset{z}{\text{const}} \left( \frac{1}{(1 - z^{c_1})(1 - z^{c_2}) \cdots (1 - z^{c_d}) z^{tb}} \right) x^t \\ &= \underset{z}{\text{const}} \left( \frac{1}{(1 - z^{c_1})(1 - z^{c_2}) \cdots (1 - z^{c_d})} \sum_{t \geq 0} \frac{x^t}{z^{tb}} \right) \\ &= \underset{z}{\text{const}} \left( \frac{1}{(1 - z^{c_1})(1 - z^{c_2}) \cdots (1 - z^{c_d})} \frac{1}{1 - \frac{x}{z^b}} \right) \quad \square \end{aligned}$$

- ① The language of polytopes
- ② The unit cube
- ③ The standard simplex
- ④ The Bernoulli polynomials as lattice-point enumerators of pyramids
- ⑤ The lattice-point enumerators of the cross-polytopes
- ⑥ Pick's theorem
- ⑦ Polygons with rational vertices
- ⑧ Euler's generating function for general rational polytopes

# Summary

- Definition of a polytope, and related concepts
- Observation of common phenomena through various examples
  - Lattice-point enumerators are polynomials in  $t$
  - Evaluation at  $-t$  gives the lattice-point enumerator of the interior
- Definition of a quasipolynomial
- Lattice-point enumerators of rational polytopes