Nash equilibrium

 $\begin{array}{l} \underline{\text{Definition 8.D.1}}: \text{(Nash equilibrium)}\\ \text{s} = (\text{s}_1, \ldots, \text{s}_{\text{I}}) \text{ is a } \underline{\text{Nash equilibrium}}\\ \text{in } \Gamma_{\text{N}} = [\text{N} = \{0, 1, \ldots, I\}, \{\text{S}_i\}, \{u_i\}]\\ \text{if } \forall i = 1, \ldots, I, \ u_i(\text{s}_i, \text{s}_{-i}) \geq u_i(\text{s}'_i, \text{s}_{-i}) \forall \text{ s}'_i \in \text{S}_i. \end{array}$

<u>Note</u>: Nash eq. → each player's strategy is a best response to the strategies actually played by her rivals Rationalizable strategies

 \rightarrow best response to some justified strategies of the rivals

Nash equilibrium

Example 8.D.1:

denotes a best response(M, m) is the unique Nash eq.

Example 8.D.2:

 (a_2, b_2) is the unique Nash eq.

rationalizable strategies \rightarrow { a_1, a_2, a_3 } for 1, { b_1, b_2, b_3 } for 2

	1	m	r
U	<u>5</u> , 3	0, 4	3, <u>5</u>
Μ	4,0	<u>5, 5</u>	4,0
D	3, <u>5</u>	0, 4	<u>5</u> , 3

	b ₁	b ₂	b ₃	b ₄
a ₁	0, <u>7</u>	2, 5	<u>7</u> , 0	0, 1
a ₂	5,2	<u>3, 3</u>	5, 2	0, 1
a ₃	<u>7</u> , 0	2, 5	0, <u>7</u>	0, 1
a ₄	0, <u>0</u>	0, -2	0, <u>0</u>	<u>10</u> , -1

<u>Note</u>: Every strategy in Nash eq. \rightarrow rationalizable

Nash equilibrium

Example 8.D.3:

_ denotes a best response

(E, E), (C, C) are Nash eq.

	E	С
Е	<u>100, 100</u>	0, 0
С	0, 0	<u>100, 100</u>

Nash eq. theory says nothing which eq. we should expect.

Best-response correspondence

$$b_{i}: S_{-i} \to S_{i}$$

$$b_{i}(s_{-i}) = \{s_{i} \in S_{i} \mid u_{i}(s_{i}, s_{-i}) \ge u_{i}(s'_{i}, s_{-i}) \forall s'_{i} \in S_{i}\}$$

$$s = (s_1, \dots, s_I) \text{ is a } \underline{\text{Nash equilibrium}}$$

in $\Gamma_N = [N = \{0, 1, \dots, I\}, \{S_i\}, \{u_i\}]$
iff $s_i \in b_i(s_{-i}) \quad \forall i = 1, \dots, I$

Nash equilibrium — Discussion

Why should we concern ourselves with the concept of Nash eq. ? How do players reach a Nash eq. ?

- 1. Nash eq. as a consequence of rational inference
- 2. Nash eq. as a necessary condition
 - if there is a unique predicted outcome
- 3. Focal points
- 4. Nash eq. as a self-enforcing agreement
- 5. Nash eq. as a stable social convention

$$\begin{array}{l} \underline{\text{Definition 8.D.1}}:\\ \sigma = (\sigma_1, \ldots, \sigma_I) \text{ is a } \underline{\text{Nash equilibrium}}\\ \text{ in } \Gamma_N = [N = \{0, 1, \ldots, I\}, \{\Delta(S_i)\}, \{u_i\}]\\ \text{ if } \forall i = 1, \ldots, I, \ u_i(\sigma_i, \sigma_{-i}) \geq u_i(\sigma'_i, \sigma_{-i}) \forall \ \sigma'_i \in \Delta(S_i). \end{array}$$

Example 8.D.4 :
((1/2, 1/2), (1/2, 1/2)) is a unique Nash eq.
1's payoff: H
$$-1 \times 1/2 + 1 \times 1/2 = 0$$

T $1 \times 1/2 + (-1) \times 1/2 = 0$
same for 2

	Н	Т
Η	-1, +1	+1, -1
Т	+1, -1	-1, +1

 $\begin{array}{ll} \underline{Proposition \ 8.D.1:} & S_{i}^{+} \subseteq S_{i} \ \text{set of pure str. played with positive prob.} \\ \text{in } \sigma = (\sigma_{1}, \ldots, \sigma_{I}). & \sigma \ \text{ is a Nash eq. in} \\ \Gamma_{N} = [N = \{0, 1, \ldots, I\}, \ \{\Delta(S_{i})\}, \ \{u_{i}\}] & \text{iff } \forall \ i = 1, \ldots, I, \\ & (i) \ u_{i}(s_{i}, \sigma_{-i}) = u_{i}(s_{i}^{*}, \sigma_{-i}) \ \forall \ s_{i}, \ s_{i}^{*} \in S_{i}^{+} \\ & (ii) \ u_{i}(s_{i}, \sigma_{-i}) \geq u_{i}(s_{i}^{*}, \sigma_{-i}) \ \forall \ s_{i} \in S_{i}^{+} \forall \ s_{i}^{*} \notin S_{i}^{+} \end{array}$

 $\begin{array}{ll} \underline{Pf}: \rightarrow) & \text{First show that } \forall \ i=1, \ \dots, I \\ & u_i(s_i, \ \sigma_{-i}) \geq u_i(s'_i, \ \sigma_{-i}) \ \forall \ s_i \in S^+_{i,} \ \underline{\forall \ s'_{\underline{i}} \in S_{\underline{i}}} \\ \text{Suppose not, i.e., } \exists \ i, \ s_i \in S^+_{i}, \ s'_i \in S_i \ s.t. \ u_i(s'_i, \ \sigma_{-i}) > u_i(s_i, \ \sigma_{-i}). \\ \text{Let } \sigma'_i \ \text{be s.t.} \end{array}$

$$\sigma'_{i}(s''_{i}) = \sigma_{i}(s''_{i}) \qquad \text{for } s''_{i} \neq s_{i}, s'_{i}$$
$$= \sigma_{i}(s'_{i}) + \sigma_{i}(s_{i}) \qquad \text{for } s''_{i} = s'_{i}$$
$$= 0 \qquad \text{for } s''_{i} = s_{i}$$

Then $u_i(\sigma'_i, \sigma_{-i}) > u_i(\sigma_i, \sigma_{-i})$, contradicting that σ is a Nash eq.

 $\begin{array}{ll} \underline{Proposition \ 8.D.1} \colon \ S^+_i \subseteq S_i \ \text{set of pure str. played with positive prob.} \\ \text{in } \sigma = (\sigma_1, \ldots, \sigma_I). \ \sigma \ \text{is a Nash eq. in} \\ \Gamma_N = [N = \{0, 1, \ldots, I\}, \ \{\Delta(S_i)\}, \ \{u_i\}] \ \text{iff } \forall \ i = 1, \ldots, I, \\ (i) \ u_i(s_i, \sigma_{-i}) = u_i(s^{\prime}_i, \sigma_{-i}) \ \forall \ s_i, \ s^{\prime}_i \in S^+_i \\ (ii) \ u_i(s_i, \sigma_{-i}) \geq u_i(s^{\prime}_i, \sigma_{-i}) \ \forall \ s_i \in S^+_i, \ \forall \ s^{\prime}_i \notin S^+_i \end{array}$

Pf: →) Next show that
$$\forall i = 1, ..., I$$

 $u_i(s_i, \sigma_{-i}) = u_i(s'_i, \sigma_{-i}) \forall s_i, s'_i \in S^+_{i,i}$

This is clear from the fact shown above:

$$u_{i}(s_{i}, \sigma_{-i}) \geq u_{i}(s'_{i}, \sigma_{-i}) \quad \forall \ s_{i} \in S^{+}_{i,} \ \underline{\forall \ s'_{\underline{i}} \in S_{\underline{i}}}$$

 $\begin{array}{ll} \underline{Proposition \ 8.D.1} \colon \ S^+_i \subseteq S_i \ \text{set of pure str. played with positive prob.} \\ \text{in } \sigma = (\sigma_1, \ldots, \sigma_I). \ \sigma \ \text{ is a Nash eq. in} \\ \Gamma_N = [N = \{0, 1, \ldots, I\}, \ \{\Delta(S_i)\}, \ \{u_i\}] \ \text{iff} \quad \forall \ i = 1, \ldots, I, \\ (i) \ u_i(s_i, \sigma_{-i}) = u_i(s^{\prime}_i, \sigma_{-i}) \ \forall \ s_i, \ s^{\prime}_i \in S^+_i \\ (ii) \ u_i(s_i, \sigma_{-i}) \geq u_i(s^{\prime}_i, \sigma_{-i}) \ \forall \ s_i \in S^+_i, \ \forall \ s^{\prime}_i \notin S^+_i \end{array}$

 $\begin{array}{ll} \underline{Pf}: \leftarrow) & \text{Suppose that } \sigma \ \text{ is not a Nash eq. }.\\ \\ \text{Then } \exists \ i, \ \sigma'_i \in \Delta(S_i) \ \text{ s.t. } u_i \ (\sigma'_i, \ \sigma_{-i}) > u_i \ (\sigma_i, \ \sigma_{-i}).\\ \\ \text{Then } \exists \ s'_i \in S_i \ \text{ s.t. } u_i \ (s'_i, \ \sigma_{-i}) > u_i \ (\sigma_i, \ \sigma_{-i}) \ \text{ with } \ \sigma'_i(s'_i) > 0.\\ \\ \\ \text{From (i), } u_i(s_i, \ \sigma_{-i}) = u_i(\sigma_i, \ \sigma_{-i}) \ \text{for all } s_i \in S^+_i.\\ \\ \\ \text{Thus } s'_i \notin S^+_i, \ \text{ contradicting (ii).} \end{array}$

Note: To see a Nash eq. or not,

it suffices to check deviations to pure strategies.

Corollary 8.D.1:

s = (s₁, ..., s_I) is a Nash eq. of $\Gamma_N = [N = \{0, 1, ..., I\}, \{S_i\}, \{u_i\}]$ iff it is a Nash eq. of $\Gamma'_N = [N = \{0, 1, ..., I, \} \{\Delta(S_i)\}, \{u_i\}]$

$\underline{Pf}: \leftarrow$) clear.

→) Since s is a Nash eq. of $\Gamma_N = [N = \{0, 1, ..., I\}, \{S_i\}, \{u_i\}],$ $\forall i = 1, ..., I \quad u_i(s_i, s_{-i}) \ge u_i(s'_i, s_{-i}) \quad \forall s'_i \notin S^+_i$

Thus (i), (ii) in Proposition 8.D.1 trivially hold since $S_{i}^{+} = \{s_i\}$ Thus by Prop.8.D.1,

s is a Nash eq. of $\Gamma'_{N} = [N = \{0, 1, ..., I, \} \{\Delta(S_{i})\}, \{u_{i}\}].$

Example 8.D.5:

S

S's mixed strategy: $(\sigma_s, 1-\sigma_s)$ T: play E $\rightarrow 1000\sigma_s$ play C $\rightarrow 100(1-\sigma_s)$

		E	С
Т	E	<u>1000, 1000</u>	0, 0
T	С	0, 0	<u>100, 100</u>

Suppose T's mixed strategy (σ_T , 1- σ_T) satisfies $0 < \sigma_T < 1$. Then S⁺_T={E,C}. Prop. 8.D.1 $\rightarrow 1000\sigma_s=100(1-\sigma_s)$ $\rightarrow \sigma s = 1/11 \rightarrow S$'s mixed strategy (1/11, 10/11)

Similarly, T's strategy (1/11, 10/11)

Nash eq. ((1/11, 10/11), (1/11, 10/11))

What is a mixed strategy ?

It just makes the rival indifferent over his strategies (The player has no preference over the probabilities.)

Is a mixed strategy useful ?

- 1 Players have a pure strategy that gives the same payoff.
 - \rightarrow why randomize them ?
 - → Players may not actually randomize; but they make definite choices that are affected by signals.
- 2 Stability of mixed strategy Nash eq.players do not have an incentive to use the exact probability
 - \rightarrow may not arise as a social convention, but as a self-enforcing agreement

Correlated Strategies



Example 8.D.5:

Public signal $\theta \in [0,1]$

- $\theta \ge 1/2 \rightarrow \text{both play E}$
- $\theta < 1/2 \ \ \rightarrow \ \ both \ \, play \ C$

		Е	С
Т	E	<u>1000, 1000</u>	0, 0
	С	0, 0	<u>100, 100</u>

This is equilibrium.

If T (S) follows, then S (T) has no incentive to deviate.

Correlated equilibrium

Proposition 8.D.2:

 $\Gamma_N = [N = \{0, 1, ..., I\}, \{\Delta(S_i)\}, \{u_i\}]$ in which $S_1, ..., S_I$ have a finite number of elements has a mixed strategy Nash eq.

Proposition 8.D.3:

A Nash eq. exists in $\Gamma_N = [N=\{0,1,\ldots,I\}, \{S_i\}, \{u_i\}]$ if $\forall i=1, \ldots, I$ (i) S_i is a nonempty, convex, and compact subset of \Re^m , and (ii) $u_i(s_1, \ldots, s_I)$ is continuous in (s_1, \ldots, s_I) and quasi-concave in s_i .

$$\begin{aligned} u_{i}(s_{1}, \dots, s_{I}) \text{ is } \underline{\text{quasi-concave}} \text{ in } s_{i} \\ \text{ if } \forall s'_{i}, s''_{i}, \alpha \in [0, 1] \\ u_{i}(\alpha s'_{i} + (1 - \alpha) s''_{i}, s_{-i}) \geq \min(u_{i}(s'_{i}, s_{-i}), u_{i}(s''_{i}, s_{-i})) \end{aligned}$$

Assignments

Problem Set 4 (due May 12):

Exercises (pp.262-266):

- 1. 8.D.3, 8.D.4, 8.D.5, 8.D.9
- Read (i) (v) on the concept of Nash equilibrium (pp.248-249) and summarize them.

Reading Assignments:

Text Chapter 8, pp.253-257