

# Trembling-Hand Perfect Equilibrium

Figure 8.F.1

		2	
1		L	R
	U	<u>1</u> , <u>1</u>	<u>0</u> , -3
	D	-3, <u>0</u>	<u>0</u> , <u>0</u>

Player 1: U w-dom D, Player 2: L w-dom R

→ (D, R) is a Nash eq. ???

((U, L) is also a Nash eq.)

## Perturbed Game

$\Gamma_\varepsilon = [N=\{0,1,\dots,I\}, \{\Delta_\varepsilon S_i\}, \{u_i\}]$  is a perturbed game of

$\Gamma_N = [N=\{0,1,\dots,I\}, \{\Delta S_i\}, \{u_i\}]$  if

$\forall i \in N, \forall s_i \in S_i \quad \exists \varepsilon_i(s_i) \in (0, 1)$  with  $\sum_{s_i \in S_i} \varepsilon_i(s_i) < 1$  s.t.

$\Delta_\varepsilon(S_i) = \{\sigma_i \mid \sigma_i(s_i) \geq \varepsilon_i(s_i) \quad \forall s_i \in S_i \text{ and } \sum_{s_i \in S_i} \sigma_i(s_i) = 1\}$

Definition 8.F.1: A Nash eq.  $\sigma$  of  $\Gamma_N = [N=\{0,1,\dots,I\}, \{\Delta S_i\}, \{u_i\}]$

is trembling-hand perfect if  $\exists$  a sequence of perturbed games

$\{\Gamma_{\varepsilon^k}\}_{k=1}^\infty$  converging to  $\Gamma_N$  (i.e.,  $\varepsilon_i^k(s_i) \rightarrow 0$  for all  $i$  and  $s_i \in S_i$

for which  $\exists$  some sequence of Nash eq.  $\{\sigma^k\}_{k=1}^\infty$  that converges to  $\sigma$ .

# Trembling-Hand Perfect Nash Equilibrium

Proposition 8.F.1: A Nash eq. of  $\Gamma_N = [N=\{0,1,\dots,I\}, \{\Delta S_i\}, \{u_i\}]$  is trembling-hand perfect iff  $\exists$  a sequence of totally mixed strategies  $\{\sigma^k\}_{k=1}^\infty$  such that  $\lim_{k \rightarrow \infty} \sigma^k = \sigma$  and  $\sigma_i$  is a best response to every element of sequence  $\{\sigma_{-i}^k\}_{k=1}^\infty$  for all  $i = 1, \dots, I$ .

Totally mixed strategy:

every pure strategy is played with positive probability

Proposition 8.F.2: If  $\sigma = (\sigma_1, \dots, \sigma_I)$  is a trembling-hand perfect Nash eq., then  $\sigma_i$  is not a weakly dominated strategy for any  $i = 1, \dots, I$ . Hence, in any trembling-hand perfect Nash eq., no weakly dominated pure strategy can be played with positive probability.

# Trembling-Hand Perfect Nash Equilibrium

Proposition 8.F.2: If  $\sigma = (\sigma_1, \dots, \sigma_I)$  is a trembling-hand perfect Nash eq., then  $\sigma_i$  is not a weakly dominated strategy for any  $i = 1, \dots, I$ . Hence, in any trembling-hand perfect Nash eq., no weakly dominated pure strategy can be played with positive probability.

$\sigma = (\sigma_1, \dots, \sigma_I)$  is a T-HPNE  $\rightarrow \sigma_i$  is not weakly dominated

Any NE not having a weakly dominated strategy  $\rightarrow$  T-HPNE ?

true for two-person games; not true in general

Existence of T-HPNE:

Every game  $\Gamma_N = [N = \{0, 1, \dots, I\}, \{\Delta S_i\}, \{u_i\}]$  with finite  $S_1, \dots, S_I$  has a T-HPNE.

## Existence of Nash Equilibrium

Lemma 8.AA.1: If  $S_1, \dots, S_I$  are nonempty, compact and convex, and  $u_i$  is continuous in  $(s_1, \dots, s_I)$  and quasi-concave in  $s_i$ , then player  $i$ 's best-response correspondence  $b_i$  is nonempty, convex-valued, and upper hemi-continuous.

Pf:  $b_i(s_{-i}) = \{s_i \in S_i \mid u_i(s_i, s_{-i}) = \max \{u_i(s'_i, s_{-i}) \mid s'_i \in S_i\}$

Non-emptiness:  $S_i$  is compact and  $u_i$  is continuous; so  $b_i(s_{-i})$  is nonempty.

Convex-valued: Pick any  $s_i, t_i \in b_i(s_{-i})$  and any  $\alpha \in [0, 1]$ . Then

$$u_i(s_i, s_{-i}) = u_i(t_i, s_{-i}) \geq u_i(s'_i, s_{-i}) \quad \forall s'_i \in S_i.$$

By the quasi-concavity of  $u_i$ ,

$$u_i(\alpha s_i + (1 - \alpha)t_i, s_{-i}) \geq \min(u_i(s_i, s_{-i}), u_i(t_i, s_{-i})) \geq u_i(s'_i, s_{-i}) \quad \forall s'_i \in S_i$$

## Existence of Nash Equilibrium

Lemma 8.AA.1: If  $S_1, \dots, S_I$  are nonempty, compact and convex, and  $u_i$  is continuous in  $(s_1, \dots, s_I)$  and quasi-concave in  $s_i$ , then player  $i$ 's best-response correspondence  $b_i$  is nonempty, convex-valued, and upper hemi-continuous.

Pf:  $b_i(s_{-i}) = \{s_i \in S_i \mid u_i(s_i, s_{-i}) = \max \{u_i(s'_i, s_{-i}) \mid s'_i \in S_i\}$

uhc: Suffice to show that for any sequence  $(s_i^n, s_{-i}^n) \rightarrow (s_i, s_{-i})$  with  $s_i^n \in b_i(s_{-i}^n) \forall n=1,2,\dots$ ,  $s_i \in b_i(s_{-i})$ .

Since  $s_i^n \in b_i(s_{-i}^n)$ ,  $u_i(s_i^n, s_{-i}^n) \geq u_i(s'_i, s_{-i}^n) \forall s'_i \in S_i$ . Thus by the continuity of  $u_i$ , we have  $u_i(s_i, s_{-i}) \geq u_i(s'_i, s_{-i}) \forall s'_i \in S_i$ .

# Existence of Nash Equilibrium

Proposition 8.D.3: A Nash equilibrium of

$\Gamma_N = [N=\{0,1,\dots,I\}, \{S_i\}, \{u_i\}]$  exists if for all  $i = 1, \dots, I$ ,

- (i)  $S_i$  is a nonempty, convex, and compact subset of some Euclidean space  $\mathbb{R}^M$ .
- (ii)  $u_i$  is continuous in  $(s_1, \dots, s_I)$ , and quasi-concave in  $s_i$ .

Pf: Define  $b: S(=S_1 \times \dots \times S_I) \rightarrow 2^S$  by  $b(s_1, \dots, s_I) = b_1(s_{-1}) \times \dots \times b_I(s_{-I})$ .

$S$  is nonempty, convex, and compact. From Lemma 8.AA.1,

$b(s_1, \dots, s_I)$  is a nonempty, convex-valued, and uhc correspondence.

Hence by the Kakutani fixed point theorem, there exists  $s \in S$

such that  $s \in b(s)$ . Therefore  $s_i \in b_i(s_{-i}) \forall i = 1, \dots, I$  which shows that

$(s_1, \dots, s_I)$  is a Nash eq.

## Existence of Nash Equilibrium

Proposition 8.D.2: Every game  $\Gamma_N = [N = \{1, \dots, I\}, \{\Delta(S_i)\}, \{u_i\}]$  in which  $S_1, \dots, S_I$  are finite sets has a mixed strategy Nash eq.

Pf:  $\Delta(S_i)$  and expected payoff functions satisfy the assumptions of Proposition 8.D.3.



# Assignments

Problem Set 6 (due May 26)

Exercises (pp.262-266): 8.F.2

Reading Assignment:

Text, Chapter 9, pp.267-276