


Nuclear Reactor Physics Lecture Note (1)  
 -Nuclear Reactions and Nuclear Cross Sections-

Toru Obara  
 Tokyo Institute of Technology


## 1. Nuclear Reactions and Nuclear Cross Sections

### 1.1 Reactions between Neutrons and nuclei


- Nuclear fission (n, fission) :



- Radioactive capture (n,  $\gamma$ ) : excited state



- Scattering (n,n) or (n,n') :



### 1.2 Microscopic Cross Sections

The probability that a neutron-nuclear reaction will occur is characterized by a quantity called a nuclear cross section.

#### (1) Definition

Consider a beam of neutrons travelling with the same speed and direction and a sufficiently thin target (one atomic layer thick).

The rate R at which reactions occur per unit area on the target:

$$R = \sigma \cdot I \cdot N_A \quad [\text{cm}^{-2} \cdot \text{s}^{-1}]$$


The microscopic cross section  $\sigma$  is defined as

$$\begin{aligned} \sigma &= \frac{(R/N_A)}{I} [\text{cm}^2] \\ &= \frac{(\text{Number of reactions/nucleus/sec})}{(\text{Number of incident neutrons/cm}^2/\text{sec})} \end{aligned}$$

The unit : b (barn)

$$1\text{b} = 10^{-24}\text{cm}^2$$

## (2) Type of neutron cross sections



Microscopic cross sections depend on the nucleus, reactions and incident neutron speed (energy)

## 1.3 Macroscopic Cross Sections

### (1) Definition

Consider a monoenergetic neutron beam incident on the surface of a target of arbitrary thickness.

The total reaction rate per unit area in  $dx$ :

$$dR = \sigma_t I N dx$$

The  $dR$  can be equal to the decrease in beam intensity between  $x$  and  $x+dx$  (with the prescription that any type of interactions will remove an incident neutron)

$$-dI(x) = \sigma_t I N dx$$

$$\therefore \frac{dI}{dx} = -N\sigma_t I(x)$$

$$\therefore I(x) = I_0 \exp(-\sigma_t N x)$$

The definition of total macroscopic cross section

$$\Sigma_t \equiv \sigma_t N [\text{cm}^{-1}]$$

## (2) Neutron mean free path and collision frequency

(Probability that a neutron moves a distance  $x$  without any interaction)

$$= \frac{I_0 \exp(-\Sigma_t x)}{I_0} = \exp(-\Sigma_t x)$$

So

(probability that a neutron has its first interaction in  $dx$ )

$$= \exp(-\Sigma_t x) \cdot \Sigma_t dx \equiv p(x) dx$$

We can calculate the average distance a neutron travels before interacting with a nucleus in the sample.

$$\begin{aligned} \bar{x} &\equiv \int_0^\infty x p(x) dx = \Sigma_t \int_0^\infty x \exp(-\Sigma_t x) dx \\ &= \frac{1}{\Sigma_t} \quad (\text{neutron mean free path}) \end{aligned}$$

$\Sigma_t$  is the probability per unit path length that a neutron will undergo a reaction, so

$$\begin{aligned} (\text{The collision frequency}) &= v \Sigma_t [\text{s}^{-1}] \\ &\text{where } v \text{ is the neutron speed.} \end{aligned}$$

## (3) Macroscopic cross sections for specific reaction

ex. macroscopic fission cross section

$$\Sigma_f = \sigma_f N$$

macroscopic absorption cross section

$$\Sigma_a = \sigma_a N$$

## (4) Macroscopic cross sections for mixture

ex. The total cross section of homogenized mixture of three different species of nuclide X, Y, and Z:

$$\Sigma_t = N_X \sigma_t^X + N_Y \sigma_t^Y + N_Z \sigma_t^Z$$