

Lecture 4 (Corrected Version)

4 Geometric Constructuion

4.1 Naive Thought in Euclidean Age

- What is a geometric construction ?
- Tools : **Compass** and **Ruler** without marking.
- **Example** : How to draw a bisector for a given angle !
- **Question** : Is it possible to construct a trisector for a give angle ?

4.2 Formulate Problem

- Start with two points on the plane, say $(0,0)$ and $(1,0)$ in euclidean coordinates. What we can do is
 1. To draw a line through two exsiting points on the plane.
 2. To record the distance between two exsiting points by compass.
 3. To draw a circle on the plane where the center is an existing point and a radius is a recorded distance.
 4. To record the interseccion of two curves (line and/or circle) on the plane.
- **Exercise** :
 1. Draw an equilatelal tiangle !
 2. Draw a square !
 3. Draw a regular pentagon (Homework 3)!
- **Exercise** :
 1. If $a, b \in \mathbb{R}$ are recorded, then $a \pm b$ can be recorded.
 2. If $a, b \in \mathbb{R}$ are recorded, then ab can be recorded.
 3. If $a, b \in \mathbb{R}$ are recorded, then a/b can be recorded.
 4. If $a \in \mathbb{R}$ is recorded, then \sqrt{a} can be recorded.

- Again start with two initial points $p_0 = (0, 0)$ and $p_1 = (1, 0)$ on the plane. Suppose we recorded m points P_m on the plane by the construction above, say,

$$P_m = \{p_0, p_1, \dots, p_m \in \mathbb{R}^2\}.$$

Using the coordinates of $p_i = (x_i, y_i)$, arrange the numbers such as

$$x_0, y_0, x_1, y_1, x_2, y_2, \dots.$$

We then let Q_n be the set consisting of the first n distinct numbers in the above ordered sequence. Thus, we obtained an increasing sequence

$$\{0, 1\} = Q_2 \subset Q_3 \subset \dots \subset Q_n \subset \dots$$

of the sets of recorded real numbers.

- Letting K_n be the field generated by Q_n over \mathbb{Q} , and we obtain a sequence of simple field extensions :

$$\mathbb{Q} = K_2 \subset K_3 \subset \dots \subset K_n \subset \dots$$

- **Theorem 4.1.1** : $[K_{n+1} : K_n] = 1$ or 2 .

Proof. The equations of a line and a circle on the plane are

$$\begin{aligned} ax + by &= c, \\ (x - d)^2 + (y - e)^2 &= r^2 \end{aligned}$$

in general. If these are drawn in terms of points in P_m , then a, b, c, d, e, r^2 can be chosen from K_n for some n . Each coordinate of an upcoming point is a root u of a quadratic equation over K_n . If $u \in K_n$, then $[K_{n+1} : K_n] = 1$. If $u \notin K_n$, then $[K_{n+1} : K_n] = 2$. \square

- **Corollary 4.1.2** : $[K_n : \mathbb{Q}] = 2^m$ for some $m \leq n$.

4.3 Three Problems in Greek Age

- **Squaring the Circle** : Construct a square with the same area as a given circle.

Proof. Impossible because π is transcendental ! \square

- **Doubling the Cube** : Construct a side of a cube that has twice the volume of a cube with a given side.

Proof. Impossible because $[\mathbb{Q}(\sqrt[3]{2}) : \mathbb{Q}] = 3$. \square

- **Trisecting the Angle** : construct an angle that is one-third of a given arbitrary angle.

Proof. Will see impossible for $2\pi/3$! \square

4.4 Homework

1. Is it always possible to construct $\alpha \in \mathbb{R}$ if $[\mathbb{Q}(\alpha) : \mathbb{Q}] = 2^n$ where $n \geq 2$?
2. Show that the equilateral n -gon can be constructed if and only if $\varphi(n) = 2^k$ for some $k \geq 1$, and if and only if $n = 2^m p_1 p_2 \cdots p_k$ so that each p_i is a distinct prime of the form $2^{2^s} + 1$ (called Fermat prime).
3. Construct the regular pentagon.
4. Construct the regular 17-gon.
5. Is it possible to construct a square from a triangle with the same area ?
6. Find an angle for which the construction of trisector is possible with respect to the initial set $P_1 = \{(0, 0), (1, 0)\}$.