



### (10) Correspondence to the free electron model

$$E = \alpha + 2\beta \cos ka$$




Upper half of the band is approximated by free electron with  $m < 0$ .  
Opposite response to electric field,  
 $\rightarrow$  **Hole** with + charge

### (11) Electron band



### (12) 2-Dimensional Square Lattice


$$\phi = \sum_n \sum_m e^{inka} e^{imka} \chi_{nm}$$



$$E \propto \sum_n \sum_m \sum_p \sum_q e^{i(n-m)ka} e^{i(p-q)ka} \int \chi_m^* H \chi_n d\tau$$


$$E = e^{ika} \beta + \alpha + e^{-ika} \beta + e^{ika} \beta + e^{-ika} \beta$$

$$E = \alpha + 2\beta \cos k_x a + 2\beta \cos k_y a$$



### (12) 2-Dimensional Square Lattice

$$E = \alpha + 2\beta \cos k_x a + 2\beta \cos k_y a$$




Bandwidth:  $8\beta$

(In general, bandwidth is  $2z\beta$ , where  $z$  is the coordination number (the number of the nearby sites.))





### (12) 2-Dimensional Square Lattice

$$E = \alpha + 2\beta \cos k_x a + 2\beta \cos k_y a$$



### (12) 2-Dimensional Square Lattice


Copper oxide high-temperature superconductor



### (13) Tight-binding method for more than two atoms in a cell

LCAO-MO  $\phi = \sum_i c_i \chi_i$

For 2 atoms,  $i=1, 2$



Bloch function is made from  $\chi_i$  for the whole crystal.

$$\chi_i \rightarrow \sum_n e^{inka} \chi_i(n)$$

Crystal orbital is constructed instead of molecular orbital as,

$$\phi = \sum_i \sum_n c_i e^{inka} \chi_i(n)$$

Calculate  $E = \frac{\int \phi^* H \phi d\tau}{\int \phi^* \phi d\tau}$  from this, and similarly to the usual Mo theory,

$\frac{\partial E}{\partial c_i} = 0$  leads to simultaneous equations of  $c_i$ , and the secular equation is

$$\begin{vmatrix} \alpha_{11}(k) - E & \beta_{12}(k) & \dots \\ \beta_{21}(k) & \alpha_{22}(k) - E & \dots \\ \vdots & \vdots & \ddots \end{vmatrix} = 0$$

Since  $\chi_i$  is Bloch function, the matrix elements,  $\alpha$  and  $\beta$  are functions of  $k$ :

$$\alpha_{ii}(k) = \int (\sum_m e^{-imka} \chi_i^*(m)) H (\sum_n e^{inka} \chi_i^*(n)) d\tau$$

$$= \alpha_i + \sum_n \beta_{ii}(n) e^{inka} \quad \beta_{ii}(n) = \int \chi_i^*(0) H \chi_i(n) d\tau$$

$$\beta_{ij}(k) = \int (\sum_m e^{-imka} \chi_i^*(m)) H (\sum_n e^{inka} \chi_j^*(n)) d\tau$$

Nearby atoms

$$= \sum_n \beta_{ij}(n) e^{inka}$$


When interaction  $\beta$  exists in the  $r$  direction, add a term  $\beta e^{ikr}$ .

$$\beta_{ij}(n) = \int \chi_i^*(0) H \chi_j(n) d\tau$$

Nearby atoms

# (14) Energy band of one-dimensional alternating chain

$\beta$  is alternately  $\beta_1$  and  $\beta_2$ .  
→ Two atoms 1 and 2 in a cell.



For simplicity, put  $\langle \chi_1 | H | \chi_1 \rangle = \langle \chi_2 | H | \chi_2 \rangle = \alpha = 0$ .

$$\langle \chi_1 | H | \chi_2 \rangle = \langle \chi_2 | H | \chi_1 \rangle^* = \beta_2 e^{-ika} + \beta_1 e^{ika}$$

$\beta_2$  in the  $-a$  direction from 1


$\beta_1$  in the  $a$  direction from 1

Secular equation is,


$$\begin{vmatrix} -E & \beta_2 e^{-ika} + \beta_1 e^{ika} \\ \beta_2 e^{ika} + \beta_1 e^{-ika} & -E \end{vmatrix} = 0$$

The solution is depicted in the right:






$$1 + \cos 2x = 2 \cos^2 x$$



Different  $\beta$  for single and double bonds



**Peierls Insulator:** Long periodicity generates a new energy gap and makes the system insulating.



## Energy band for a system with many atomic orbitals

LCAO-MO from all atomic orbitals in a cell  
( $N$ , each atom may have more than one.)

$$\phi = \sum_i c_i \chi_i$$



Make the Bloch function of each  $\chi$





$\chi_i \rightarrow \sum_n e^{ink_a} \chi_i(n)$  leading to an  $N \times N$  secular equation

$$\begin{vmatrix} \alpha_{11}(k) - E & \beta_{12}(k) & \dots \\ \beta_{21}(k) & \alpha_{22}(k) - E & \dots \\ \vdots & \vdots & \ddots \end{vmatrix} = 0$$


Since each element is a function of  $k$ , this secular equation is solved at each  $k$ , to give  $N$  energy levels.

These energy levels for different  $k$  are connected to afford continuous energy bands (right).







How to make the Brillouin zone.



How to make the Brillouin zone.




How to make the Brillouin zone for a oblique lattice.



Brillouin zone and  
the Bravais lattices

Reciprocal of centered  
is centered






## Brillouin zone and the Bravais lattices

Reciprocal of face centered is body centered


Reciprocal of body centered is face centered



## Brillouin zone and the Bravais lattices


体心立方格子  
bcc  
body centered  
cubic

面心立方格子  
fcc  
face centered  
cubic



## Fermi surface of metals

(1) Alkali metals Li, Na, K




Fermi surface  $\Leftrightarrow$  trajectory of  $k_F \Leftrightarrow$  Surface at  $E=E_F=\text{const.}$

Assuming free electron

$$E = \frac{\hbar^2}{2m}(k_x^2 + k_y^2 + k_z^2) = \text{const.}$$

leads to a sphere in the  $k$ -space.





Fermi surface of alkali metals is not deviated from the perfect sphere by 1%.

Energy interval in the  $k$ -space is  $\Delta k = \frac{2\pi}{Na}$


Volume of the Fermi sphere is half of the volume of the 1st Brillouin zone.

## Fermi surface of metals


(1) Divalent metals Mg, Al




Volume of the Fermi sphere is the same as the volume of the 1st Brillouin zone.



第1ゾーン







第2ゾーン




第3ゾーン

## Fermi surface of metals


### (2) Divalent and trivalent Mg, Al



第1ゾーン



第2ゾーン




第3ゾーン

## Fermi surface of metals




### (3) Transition metals

3d band is narrower than the 4s band.



## Fermi surface of metals



### (4) Cu, Ag, Au $s^1 d^{10} \rightarrow$ close to alkali metals



## 1D metal

$$E = \frac{\hbar^2 k_x^2}{2m} = \text{const.} \quad \text{leads to} \quad k_x = k_F = \text{const.}$$


(No momentum for other  $k_y$  and  $k_z$  directions.  
= cannot move.)



The Fermi surface consists of a pair of planes.


## Examples of 1D metals

### (1) 1D Platinum Complex KCP $K_2[Pt(CN)_4]Br_{0.40}xH_2O$





Pt surrounded by CN  
Metal 1D Chain

Metal at room temp.  
Insulator at low temp.





## Examples of 1D metals

### (2) NbSe<sub>3</sub>




### (3) Polyacetylene



## Examples of 1D metals



### (4) Organic Charge-Transfer Complex (TTF)(TCNQ)



Current flow

## Examples of 2D metals

$$E = \frac{\hbar^2}{2m} (k_x^2 + k_y^2) = \text{const.} \quad \text{leads to} \quad k_x^2 + k_y^2 = k_F^2 = \text{const.} \rightarrow \text{circle}$$




Cylindrical Fermi surface


Examples  
Graphite  
Organic superconductors



## Organics : energy levels → energy bands



## Fermi surface of the first organic superconductor



## (TMTSF)<sub>2</sub>PF<sub>6</sub>


$$E(k) = 2\beta_b \cos(kb) \pm \sqrt{\Delta}$$

$$\Delta = [(\beta_{a1} + \beta_{a2}) \cos(\frac{ka}{2}) + (\beta_{p1} + \beta_{p2}) \cos(\frac{ka}{2} - kb)]^2$$

$$+ [(\beta_{a1} - \beta_{a2}) \sin(\frac{ka}{2}) + (\beta_{p1} - \beta_{p2}) \sin(\frac{ka}{2} - kb)]^2$$

方向	β (meV)
HOMO	
a1	200
a2	230
b	35
p1	20
p2	7


Calculated from MO



A pair of waving planes → Quasi-1D conductor

Considerable waving reflects the 2D character.

## Fermi surface of organic superconductor




$$\beta_{11} = \beta_{22} = \beta_c e^{ikc} + \beta_c e^{-ikc}$$

$$= 2\beta_c \cos(kc)$$

$$\beta_{12} = \beta_{21}^* = \beta_{p1} + \beta_{p2} e^{i(kb+kc)}$$

$$+ \beta_{q2} e^{ikc} + \beta_{q1} e^{ikb}$$




Cylindrical Fermi surface indicated 2D.


Program available from <http://www.op.titech.ac.jp/lab/mori/lib/program.html>

## Excise Band structure of the $\theta$ -phase

This generally found structure is called herringbone structure in organic crystals or alternatively  $\theta$ -phase in organic conductors. Calculate the energy bands of this structure.



A unit cell contains two molecules, numbered 1 and 2. Transfer  $t_a$ , running along  $a$ , is between two Molecule 1. Transfer  $t_p$ , running diagonal ( $a/2, b/2$ ) etc, is between Molecule 1 and 2.



(1) How many Molecule 1 exist near Molecule 1?

From this, obtain the diagonal element  $F_{11}$ .

(1) How many Molecule 2 exist near Molecule 1?

From this, obtain the nondiagonal element  $F_{12}$ .

(3) Solve the secular equation, and obtain an equation of  $E(k_a, k_b)$  representing the energy band.

$$\cos x + \cos y = 2 \cos \frac{x+y}{2} \cos \frac{x-y}{2}$$